

02/10/98

Box Seq.

A

+

Please type a plus sign (+) inside this box → ☐

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PTO/SB/05 (12/97)

Approved for use through 09/30/00. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

**UTILITY  
PATENT APPLICATION  
TRANSMITTAL**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 6048.US.P1

Total Pages 141

First Named Inventor or Application Identifier

Steven R. Wiley

Express Mail Label No. EH 824 367 096 US

**APPLICATION ELEMENTS**

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

1. ☒ Fee Transmittal Form  
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 140]  
(preferred arrangement set forth below)
- Descriptive title of the invention
  - Cross References to Related Applications
  - Statement Regarding Fed sponsored R & D
  - Reference to Microfiche Appendix
  - Background of the invention
  - Brief Summary of the invention
  - Brief Description of the Drawings (if filed)
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure
3. ☒ Drawing(s) (35 USC 113) [Total Sheets 1]
4. Oath or Declaration [Total Pages ]
- a. ☒ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 CFR 1.63(d))  
(for continuation/divisional with Box 17 completed)  
(Note Box 5 below)
- c. ☐ DELETION OF INVENTOR(S)  
Signed statement attached deleting  
inventor(s) named in the prior application,  
see 37 CFR 1.63(d)(2) and 1.33(b).
5. ☐ Incorporation By Reference (useable if Box 4b is checked)  
The entire disclosure of the prior application, from which a  
copy of the oath or declaration is supplied under Box 4b,  
is considered as being part of the disclosure of the  
accompanying application and is hereby incorporated by  
reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. Nucleotide and/or Amino Acid Sequence Submission  
(if applicable, all necessary)
- a. ☒ Computer Readable Copy
- b. ☒ Paper Copy (identical to computer copy)
- c. ☒ Statement verifying identity of above copies

**ACCOMPANYING APPLICATION PARTS**

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement [ ] Power of Attorney  
(when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure [ ] Copies of IDS  
Statement (IDS)/PTO-1449 Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)
14. ☐ Small Entity [ ] Statement filed in prior application.  
Statement(s) Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
16. ☒ Other: Executed, Assignment  
Recordation Form Cover Sheet

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☒ Continuation-in-part (CIP) of prior application No: 08 / 798,692**18. CORRESPONDENCE ADDRESS**☐ Customer Number or Bar Code Labelor ☐ Correspondence address below

(Insert Customer No. or Attach bar code label here)

NAME	ABBOTT LABORATORIES				
	Steven F. Weinstock				
ADDRESS	Department 377/AP6D-2				
	100 Abbott Park Road				
CITY	Abbott Park	STATE	Illinois	ZIP CODE	60070-3500
COUNTRY	USA	TELEPHONE	847-935-1729	FAX	847-938-2623

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

MEMBER OF THE TNF FAMILY USEFUL FOR TREATMENT AND DIAGNOSIS  
OF DISEASE

5 The present application is a continuation-in-part of  
pending U.S. patent application serial no. 08/798,692 filed  
on February 12, 1997 which is hereby incorporated in its  
entirety by reference.

Background of the Invention

10 This invention relates generally to extracellular  
signal molecules, and more particularly, relates to a member  
of the tumor necrosis factor (TNF) family of molecules  
designated as TREPA, reagents and methods for its detection  
as well as its use in therapeutics.

15 The TNF (tumor necrosis factor) family is an expanding  
set of extracellular signaling molecules (ligands) with  
biological activities that are intimately associated with a  
variety of disease conditions. The prototypic member of  
this family, TNF, is well known as a mediator of septic  
20 shock, inflammation, and graft versus host disease. See,  
for example, A. Cerami, Immunol Today, 9:28-31 (1988); M.  
Revel, Ciba Found Symp, 129:223-33 (1987); J. Cohen, J. Bone  
Marrow Transplant 3(3):193-197 (1988). Also, because of its  
beneficial effects on vasculature of solid tumors, isolated  
25 perfusion of TNF is being evaluated as a therapeutic agent  
for cancer patients. M. W. Boehme, Eur. J. Clin. Invest.  
26: 404-410 (1996).

The important role of the TNF family in immune  
regulation has been demonstrated by mutations both in mice  
30 and humans. For instance, mice with a loss of function  
mutation in the TNF family member known as Fas ligand  
present a variety of disorders including lymphadenopathy and  
autoimmune disease. See, for example, R. Watanabe-  
Fukunaga, Nature 356:314-317 (1992); Takahishi et al., Cell  
35 76:969-966, (1994), Adiachi et al, PNAS, 90:1756 (1993);  
Fisher et al, Cell 81:953-946 (1995); F. Rieux-Laucat,  
Science 268:1347 (1995). Another example of the immune  
regulation role is mutation of the TNF family member CD40

ligand in humans. Spontaneous CD40 ligand mutations in human patients result in hyper Ig-M syndrome, demonstrating the requirement of CD40 ligand for B-cell maturation and isotype switching. Also, targeted disruption of TNF family member LTa in mice results in failure to develop peripheral lymph nodes. P. De Togni, Science 264:703-707 (1994).

Another property of this family of ligands which is potentially clinically useful is the ability to selectively induce apoptosis (programmed cell death) in a variety of cancer cells, but not in most normal cells. The two known members of the TNF family which induce apoptosis in the widest variety of cell lines are TRAIL (TNF related apoptosis inducing ligand) and Fas ligand. See, for example, T. Suda et. al., Cell 75:1169-1178 (1993); Wiley et. al., Immunity 3:673-682, (1995). This property is unique to the TNF family of ligands.

Members of the TNF family of ligands can be identified by a region of amino acid conservation which is approximately restricted to the N-terminal 150 amino acids. This region forms a b-pleated sheet structure which trimerizes and interacts with the cognate receptors. Within this N-terminal domain there are isolated regions of homology which correspond to the strands of the b-pleated sheet. Therefore, the total amino acid sequence identity between TNF family members is not high, but can be recognized by those skilled in the art.

Given the crucial roles of members of this family of molecules in immune regulation, investigation into the existence and identity of other members of the TNF family is desirable. The identification and characterization of these molecules provide means to identify and treat a variety of immune disorders.

#### Summary of the Invention

The present invention provides a novel member of the TNF family of ligands, as well as cDNA encoding this novel member, and expression vectors encoding a soluble form of this novel member. Since this novel member of the TNF

family is able to induce proliferation in human umbilical endothelium cells, it is called TREPA which stands for TNF Related Endothelium Proliferative Agent.

5 A method for producing TREPA polypeptides involves expression in host cells transformed with a recombinant expression vector that contains an engineered soluble version of TREPA, as well as a cell surface expressed form of TREPA.

10 The present invention also provides a method of detecting target polynucleotides of TREPA in a test sample which comprises contacting a target polynucleotide specific for TREPA with at least one TREPA specific polynucleotide and fragment or complement thereof provided herein and detecting the presence of the target in the test sample.  
15 The polynucleotide comprises SEQUENCE ID NO 1 and fragments or complements thereof. Also, the TREPA polynucleotide may be attached to a solid phase prior to performing the assay.

20 The present invention also provides a method for amplifying 5' end cDNA of TREPA gene in a test sample, which comprises performing reverse transcription with random primers, amplifying the cDNA obtained by using other oligonucleotide primer(s) of TREPA as sense and antisense primer(s) in a first-stage PCR to obtain amplified cDNA and detecting the presence of TREPA amplicon in the test sample.  
25 Amplification can be performed by the polymerase chain reaction. Also, the test sample can be attached to a solid phase prior to performing the method. Further, the detection step can comprise utilizing a detectable label capable of generating a measurable signal. The detectable  
30 label can be attached to a solid phase.

The present invention further provides a method of detecting TREPA in a test sample suspected of containing TREPA, which comprises contacting said test sample with at least one polynucleotide as a sense primer and with at least  
35 one polynucleotide as an anti-sense primer and amplifying same to obtain a first stage reaction product; contacting said first stage reaction product with at least one of said

polynucleotides of the contacting step and a second polynucleotide, with the proviso that the second oligonucleotide is located 3' to the first oligonucleotide utilized and is of opposite sense to said first  
5 oligonucleotide, and detecting said TREPA as an indication of the presence of disease. The amplification may be performed by the polymerase chain reaction. The test sample can be attached to a solid phase prior to performing the method. The detection step also comprises utilizing a  
10 detectable label capable of generating a measurable signal, and the detectable label can be attached to a solid phase. Test kits useful for detecting TREPA target in a test sample further are provided which comprise a container containing at least one polynucleotide selected from the group  
15 consisting of SEQUENCE ID NO 1, and fragments and complements thereof. These test kits further comprising containers containing tools useful for collecting test samples such as blood, urine, saliva, and stool. Such tools include lancets and absorbent paper or cloth for collecting  
20 and stabilizing blood; swabs for collecting and stabilizing saliva; cups for collecting and stabilizing urine or stool samples. Collection materials, papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. They  
25 may also be treated with or contain preservatives, stabilizers or antimicrobial agents to help maintain the integrity of the specimens.

The present invention provides a purified polynucleotide or fragment thereof derived from the TREPA  
30 gene capable of selectively hybridizing to the genomic TREPA gene or the complement thereof. The polynucleotide is selected from the group consisting of SEQUENCE ID NO 1, and fragments and complements thereof. Further, the polynucleotide can be produced by recombinant techniques.  
35 This recombinant molecule comprises a sequence that encodes at least one epitope of TREPA and is contained within a

recombinant vector. The recombinant polynucleotide further comprises a host cell transformed with said vector.

The present invention further provides a recombinant expression system comprising an open reading frame of DNA or  
5 RNA derived from the TREPA gene wherein said open reading frame comprises the test sample with at least one TREPA specific polynucleotide or complement thereof, and detecting the presence of the target polynucleotide of TREPA in the test sample. The TREPA specific polynucleotide has at least  
10 50% identity to polynucleotide SEQUENCE ID NO 1 and fragments, analogs or complements thereof. The target polynucleotide of TREPA can be attached to a solid phase prior to performing the method.

The present invention also includes a method for  
15 detecting mRNA of TREPA in a test sample, which comprises (a) performing reverse transcription with at least one primer in order to produce cDNA; (b) amplifying the cDNA obtained from step (a) by using other oligonucleotide primer(s) of TREPA as sense and antisense primer(s) in a  
20 first-stage amplification to obtain TREPA amplicon; and (c) detecting the presence of TREPA amplicon in the test sample, wherein the oligonucleotide primers of TREPA have at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof. The method further comprises reacting  
25 the test sample with a solid phase prior to performing step (a) or step (b) or step (c). Further, the detection step can further comprise utilizing a detectable label capable of generating a measurable signal.

An additional method is provided for detecting target  
30 TREPA polynucleotide in a test sample suspected of containing the target. This method comprises (a)

contacting the target TREPA polynucleotide with at least one TREPA oligonucleotide as a sense primer and with at least one TREPA oligonucleotide as an anti-sense primer  
35 and amplifying same to obtain a first stage reaction product; (b) contacting said first stage reaction product with at least one other TREPA oligonucleotide, with the

proviso that the other TREPA oligonucleotide is located 3' to the TREPA oligonucleotides utilized in step (a) and is complementary to the first stage reaction product; and (c) detecting the target TREPA polynucleotide, wherein the

5 TREPA oligonucleotides utilized in step (a) and step (b) have at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof. Further, the test sample can be reacted with a solid phase prior to performing step (a) or step (b) or step (c). Also, the

10 detection step can comprise utilizing a detectable label capable of generating a measurable signal. Further, the detectable label can be attached to a solid phase.

The present invention further provides a test kit useful for detecting TREPA polynucleotide in a test sample,

15 comprising a container containing at least one TREPA polynucleotide having at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof. Further, the test kit comprises a container containing tools useful for collection of said sample selected from the group

20 consisting lancets, absorbent paper, cloth, swabs and cups.

A purified polynucleotide or fragment thereof derived from the TREPA gene also is provided. The purified polynucleotide is capable of selectively hybridizing to the nucleic acid of said TREPA gene, and the purified

25 polynucleotide has at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof. Further, the purified polynucleotide can be produced by recombinant techniques. When produced by recombinant techniques, the purified polynucleotide can further comprise a sequence of

30 at least one epitope encoded by TREPA.

The present invention provides a recombinant expression system comprising a nucleic acid sequence that encodes an open reading frame derived from TREPA which is operably linked to a control sequence compatible with a desired host.

35 The nucleic acid sequence has at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof. This recombinant expression system further

comprises a cell transformed with the recombinant expression system.

The present invention further provides a polypeptide encoded by TREPA. The polypeptide has at least 35 %  
5 identity to amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof. The polypeptide can be produced by recombinant technology or by synthetic techniques.

A compound which inhibits activation of the TREPA  
10 polypeptide is provided. The TREPA polypeptide has at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof. The invention further provides a polypeptide as a soluble fragment of the TREPA protein which  
15 is capable of binding a receptor for TREPA.

A method for treating a patient having a need to induce activation of the TREPA polypeptide is provided, which comprises administering to the patient a therapeutically effective amount of a compound which induces activation of  
20 the TREPA polypeptide. TREPA polypeptide has at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

The present invention provides a method for determining  
25 whether a compound is an agonist or antagonist to TREPA protein. This method comprises (a) contacting a cell having TREPA protein expressed on its surface with said compound and a receptor ligand; (b) determining whether a biological effect is produced from the interaction of the ligand and  
30 the receptor; and (c) determining whether the compound is an agonist or antagonist. The protein has at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

35 The present invention further provides a method for determining whether a receptor binds to a TREPA ligand. This method comprises (a) contacting a mammalian cell which



expresses the TREPA ligand with a receptor; (b) detecting the presence of the receptor; and (c) determining whether the receptor binds to the TREPA ligand.

5 An antibody which specifically binds to at least one epitope encoded by TREPA is provided herein. The antibody is polyclonal or monoclonal. The epitope comprises an amino acid sequence having at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

10 Also provided is an assay kit for determining the presence of TREPA antigen or antibody in a test sample, which comprises a container containing a TREPA polypeptide having at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2,  
15 SEQUENCE ID NO 3 and fragments thereof. Further, the polypeptide can be attached to a solid phase. In addition, the assay kit further comprises a container containing tools useful for collection of said sample selected from the group consisting lancets, absorbent paper, cloth, swabs and cups.  
20 Another assay kit for determining the presence of TREPA antigen or antibody in a test sample which is provided by the present invention comprises a container containing an antibody which specifically binds to TREPA antigen, wherein the antigen comprises at least one epitope of TREPA having  
25 at least about 60% similarity to a sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof. Further, the assay kit comprises a container containing tools useful for collection of said sample selected from the group consisting lancets, absorbent  
30 paper, cloth, swabs and cups. Further, the antibody of the kit can be attached to a solid phase.

The present invention provides a method for producing a polypeptide comprising at least one epitope of TREPA. The method comprises incubating host cells transformed with an  
35 expression vector, wherein the vector comprises a polynucleotide sequence encoding a polypeptide, which polypeptide comprises an amino acid sequence having at least

35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

The present invention further provides a method for  
5 detecting TREPA antigen in a test sample suspected of  
containing the TREPA antigen, comprising (a) contacting the  
test sample with an antibody or fragment thereof which  
specifically binds to at least one epitope of TREPA antigen  
selected from the group consisting of SEQUENCE ID NO 2,  
10 SEQUENCE ID NO 3 and fragments thereof, for a time and under  
conditions sufficient for the formation of antibody/antigen  
complexes; (b) detecting said complexes. The method further  
comprises the antibody attached to a solid phase.

A method for detecting antibodies which bind to TREPA  
15 antigen in a test sample suspected of containing the  
antibodies is provided by the invention. The method  
comprises (a) contacting said test sample with a TREPA  
polypeptide, wherein the TREPA polypeptide contains at  
least one TREPA epitope comprising an amino acid sequence or  
20 fragment thereof having at least 35% identity to an amino  
acid sequence selected from the group consisting of SEQUENCE  
ID NO 2, SEQUENCE ID NO 3 and fragments thereof, for a time  
and under conditions sufficient to allow antigen/antibody  
complexes to form; and (b) detecting said complexes. The  
25 TREPA polypeptide further can be attached to a solid phase.

Also provided by the present invention is a tissue  
culture grown cell comprising a nucleic acid sequence that  
encodes at least one epitope of TREPA antigen or a fragment  
thereof, wherein the nucleic acid sequence is transfected  
30 into the cell and wherein the nucleic acid sequence is  
SEQUENCE ID NO 1 and fragments, analogs or complements  
thereof.

In addition, the present invention provides a method  
for producing antibodies which specifically bind to TREPA  
35 antigen. The method comprises administering to an  
individual an isolated immunogenic polypeptide or fragment  
thereof, wherein the isolated immunogenic polypeptide

comprises at least one TREPA epitope and has at least 35% identity to a sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof, in an amount sufficient to produce an immune response. An  
5 additional method for producing antibodies which specifically bind to TREPA antigen is provided, which comprises administering to a mammal a plasmid comprising a sequence which encodes at least one epitope of TREPA, wherein the TREPA sequence is selected from the group  
10 consisting of SEQUENCE ID NO 1 and fragments or complements thereof.

The present invention provides a composition of matter comprising a TREPA polynucleotide or fragment thereof, wherein the polynucleotide has at least 50% identity to  
15 SEQUENCE ID NO 1 and fragments, analogs or complements thereof. Further provided is a composition of matter comprising a polypeptide containing at least one epitope encoded by TREPA, wherein the polypeptide has at least 35% identity to a sequence selected from the group consisting of  
20 SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof. This polypeptide composition of matter further comprises a soluble fragment of the TREPA protein and is capable of binding a receptor for TREPA.

The present invention also provides a gene or fragment  
25 thereof which codes for TREPA protein which comprises an amino acid sequence which has at least 35% identity to SEQUENCE ID NO 5. In addition, the present invention provides a gene or fragment thereof comprising DNA having at least 35% identity to SEQUENCE ID NO 4.

30

#### Brief Description of the Drawings

FIGURE 1 presents an amino acid alignment of TREPA with several previously described members of the TNF family of ligands. hTREPA, or human TREPA is SEQUENCE ID NO 6, hTRAIL  
35 is SEQUENCE ID NO 7, mTRAIL is SEQUENCE ID NO 8, hTNFa is SEQUENCE ID NO 9, hTNFb is SEQUENCE ID NO 10 and h4-1BBL is SEQUENCE ID NO 11.

### Detailed Description of the Invention

A novel protein designated TREPA is provided herein, along with the DNA that encodes the receptor binding domain of TREPA, and recombinant vectors for producing TREPA DNA.

The present invention also provides antibodies that specifically bind TREPA proteins. In one embodiment, these antibodies are monoclonal antibodies.

Identification of a cDNA clone encoding human TREPA is described in Example 1 below. The nucleotide sequence of the human TREPA gene is presented in SEQUENCE ID NO 1, and the amino acid sequence of the protein is presented in SEQUENCE ID NO 2 and SEQUENCE ID NO 3.

The carboxyl-terminal ~150 amino acids of the members of the TNF family demonstrate sequence conservation. New members of the family therefore can be recognized by sequence homology to known family members in that region. The amino acid sequences disclosed herein reveal that TREPA is a member of the TNF family of ligands. Smith et al, Cell 73:1349 (1993); Suda et. al., Cell 75 :1169 (1993); Smith et. al. Cell 76:959 (1994). Of all known members of the TNF family, TREPA is most closely related to TNF-alpha.

Conserved sequences located in the carboxyl terminal portion of the TNF family of ligands are identified in Smith et. al., Cell, supra. FIGURE 1 presents an amino acid alignment of TREPA (SEQUENCE ID NO 6) with several previously described members of the TNF family of ligands. These are, from top to bottom, human TRAIL (SEQUENCE ID NO 7), murine TRAIL (SEQUENCE ID NO 8), human TNF-a (SEQUENCE ID NO 9), murine TNF-b (SEQUENCE ID NO 10), and human 4-1BB ligand (SEQUENCE ID NO 11), respectively. Several highly conserved residues that are common to a majority of TNF family member are present in TREPA. Amongst these residues are amino acids 109 (A), 110 (H), 133 (W), 159 (G), 161 (Y), 166 (Q), 169 (F), 205 (P), 213 (S), 234 (L), 243 (F), 244 (G), and 246 (F).

Therapeutic Applications

Endothelial cells are cells that line the walls of blood vessels. Proliferation of endothelial cells can result in growth of blood vessels in a tissue, a process known as vascularization. There are several clinical conditions where it is beneficial to promote vascularization of a tissue. This invention provides for use of TREPA as a therapeutic agent to promote vascularization in those situations. For example, wounded tissues require vascularization as part of the healing process. Since TREPA has been shown to increase vascularization of HUVEC cells (human umbilical vein endothelial cells) (see example XX) this invention provides for use of TREPA as a therapeutic to promote wound healing. Another situation where tissue vascularization is desirable is in tissue grafts, where the grafted tissue must be vascularized by the host in order for the graft to be successful. Therefore, the present invention provides for use of TREPA as a therapeutic agent to promote tissue grafts.

Likewise, there are situations where inhibiting vascularization is desirable. For example, formation of solid tumors requires that the tumor be vascularized by the host or the tumor cannot grow beyond a certain limited size. Inhibiting vascularization of solid tumors can therefore prevent their growth. The present invention provides for blocking of TREPA, by means such as anti-sense RNA, blocking antibody, or compounds developed by targeting TREPA in order to prevent vascularization of solid tumors.

Programmed cell death (apoptosis) is a fundamental biological process. Inflammation is a biological process whereby a region of the body is infiltrated by a variety of activated lymphocytes, in response to conditions such as bacterial infection or autoimmunity. While this process is beneficial, there are disease conditions caused by excessive and inappropriate inflammation which results in tissue damage. TNF has been implicated as a major cause of some of these conditions such as sepsis, rheumatoid arthritis and

inflammatory bowel disease. Since TREPA is closest to the TNF-alpha member of the TNF family, the present invention provides for the use of TREPA as an indicator of inflammation, and for blocking of TREPA, by means such as anti-sense RNA, blocking antibody, or compounds developed by targeting TREPA as a means to inhibit inflammation.

Programmed cell death (apoptosis) is a fundamental biological process whereby, given appropriate and external signals, cells are induced to self -destruction. This process had been described to occur during embryogenesis, endocrine-dependent tissue atrophy, normal tissue turnover, nervous and immune system development. In the case of the immune system, T-cells that recognize self epitopes are destroyed by apoptosis during maturation of T-cell in the thymus. It has been proposed that failure of self-reactive T-cells to undergo apoptosis is responsible for autoimmune disorders. Gammon et. al., Immunology Today 12: 193 (1991). One characteristic of the TNF family is the ability of many family members to induce programmed cell death (apoptosis) in a variety of cells, both normal and of tumor origin (Wiley et al, Immunity 3:673-682, (1995), and references therein) Therefore, the present invention provides for use of TREPA as an anti-cancer agent to induce apoptosis in cancer and tumor associated cells. The present invention also provides for use of TREPA as an agent to eliminate inappropriate untransformed cells such as self-reactive T-cells. TREPA may be internally administered alone or in conjunction with other agents or it may be used for the ex-vivo treatment and replacement of disease tissues. For instance circulating lymphoma cells may be exposed to immobilized TREPA ex-vivo, and the treated blood reintroduced into the patient.

Another characteristic of the TNF family of ligands is to increase exposure of solid tumors to cells of the immune system by up-regulating vascular adhesion molecules, thereby attracting anti-tumor leukocytes to the region. M. W. Boehme, Eur. J. Clin. Invest., supra. TNF also lowers the

interstitial pressure of tumors, allowing concentrations of anti-cancer chemical agents or mixtures of anti-cancer chemical agents to enter the tumor in higher concentrations. The present invention provides for use of TREPA protein as an adjunct to be used with anti-cancer chemotherapy agents for the treatment of tumors.

In addition, viral infection normal cells can make them susceptible to apoptosis induced by TNF family members, such as TNF-alpha and Fas Ligand. See, for example, C.V. Paya et. al., J. Immunol. 141:1989-1995 (1988); P.D. Katsiskas, J. Exp. Med. 181:2029-2036 (1995). The present invention provides for use of TREPA protein as an anti-viral therapeutic agent, to be used alone or in conjunction with other agents, such as anti-viral compounds, or protein therapeutic agents. Since pre-treatment of virally infected cells with g-interferon has been shown to increase apoptosis induced by members of the TNF family, one embodiment of this application is to administer TREPA protein together with other agents such as g-interferon.

This invention also provides for the use of TREPA protein in developing any disorder mediated (directly or indirectly) by insufficient amounts or production of defective TREPA protein. Purified human TREPA protein may be administered to a patient with such a condition. Alternatively, gene therapy techniques for producing TREPA peptide in vivo are also provided.

Pharmaceutical preparations of TREPA comprise purified TREPA protein and a physiologically acceptable carrier, diluent, or excipient. Such compositions include buffers, anti-oxidants, low molecular weight peptides, proteins, amino acids, carbohydrates, chelating agents, glutathione, and other excipients and stabilizers commonly found in pharmaceutical compositions.

The present invention provides methods for assaying a test sample for products of a TREPA gene, which comprise making cDNA from mRNA in the test sample, and detecting the cDNA as an indication of the presence of the TREPA gene.

The method may include an amplification step, wherein portions of the cDNA corresponding to the gene or fragment thereof are amplified. Methods also are provided for assaying for the translation products of mRNAs. Test  
5 samples which may be assayed by the methods provided herein include tissues, cells, body fluids and secretions. The present invention also provides reagents such as oligonucleotide primers and polypeptides which are useful in performing these methods.

10 Portions of the nucleic acid sequences disclosed herein are useful as primers for the reverse transcription of RNA or for the amplification of cDNA; or as probes to determine the presence of certain cDNA sequences in test samples. Also disclosed are nucleic acid sequences which permit the  
15 production of encoded polypeptide sequences which are useful as standards or reagents in diagnostic immunoassays, targets for pharmaceutical screening assays and/or as components or target sites for various therapies. Monoclonal and polyclonal antibodies directed against at least one epitope  
20 contained within these polypeptide sequences are useful for diagnostic tests as well as delivery agents for therapeutic agents and for screening for diseases or conditions associated with the TREPA gene provided herein, especially inflammation. Isolation of sequences of other portions of  
25 the gene of interest can be accomplished by utilizing probes or PCR primers derived from these nucleic acid sequences, thus allowing additional probes and polypeptides of the genome of interest to be established, which also will be useful in the diagnosis, prognosis and/or treatment of  
30 diseases and conditions characterized by the TREPA gene disclosed herein.

The techniques for determining the amino acid sequence "similarity" are well-known in the art. In general, "similarity" means the exact amino acid to amino acid  
35 comparison of two or more polypeptides at the appropriate place, where amino acids are identical or possess similar chemical and/or physical properties such as charge or



hydrophobicity. A so-termed "percent similarity" then can be determined between the compared polypeptide sequences. The techniques for determining nucleic acid and amino acid sequence identity also are well known in the art and include  
5 determining the nucleotide sequence of the mRNA for that gene (usually via a cDNA intermediate) and determining the amino acid sequence encoded therein, and comparing this to a second amino acid sequence. In general, "identity" refers to an exact nucleotide to nucleotide or amino acid to amino  
10 acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more polynucleotide sequences can be compared by determining their "percent identity." Two amino acid sequences likewise can be compared by determining their "percent identity." The  
15 programs available in the Wisconsin Sequence Analysis Package, Version 8 (available from Genetics Computer Group, Madison, WI), for example, the GAP program, are capable of calculating both the identity between two polynucleotides and the identity and similarity between two polypeptide  
20 sequences, respectively. Other programs for calculating identity or similarity between sequences are known in the art.

The compositions and methods described herein will enable the identification of certain markers as indicative  
25 of TREPA disease; the information obtained therefrom will aid in the diagnosis, staging, monitoring, prognosis and/or therapy of diseases or conditions associated with the TREPA gene, especially inflammation, cancer, and graft-vs-host disease. Test methods include, for example, probe assays  
30 which utilize the sequence(s) provided herein and which also may utilize nucleic acid amplification methods such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), and hybridization. In addition, the nucleotide sequences provided herein contain open reading frames from  
35 which an immunogenic epitope may be found. This epitope is believed to be unique to the disease state or condition associated with the TREPA gene. It is thought that the

TREPA gene described herein is useful as a marker either elevated in disease such as inflammation, altered in disease such as inflammation, or present as a normal protein but appearing in an inappropriate body compartment. The

5 uniqueness of the epitope may be determined by its immunological reactivity with the specific TREPA gene, especially diseases of inflammation, such as sepsis. Methods for determining immunological reactivity are well-known and include but are not limited to, for example,  
10 radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), hemagglutination (HA), fluorescence polarization immunoassay (FPIA), chemiluminescent immunoassay (CLIA), and others; several examples of suitable methods are described herein.

15 Unless otherwise stated, the following terms shall have the following meanings:

A polynucleotide "derived from" a designated sequence refers to a polynucleotide sequence which is comprised of a sequence of approximately at least about 6 nucleotides, is  
20 preferably at least about 8 nucleotides, is more preferably at least about 10-12 nucleotides, and even more preferably is at least about 15-20 nucleotides corresponding, i.e., identical to or complementary to, a region of the designated nucleotide sequence. The sequence may be complementary to  
25 or identical to a sequence which is unique to a particular polynucleotide sequence as determined by techniques known in the art. Comparisons to sequences in databanks, for example, can be used as a method to determine the uniqueness of a designated sequence. Regions from which sequences may  
30 be derived include but are not limited to regions encoding specific epitopes, as well as non-translated and/or non-transcribed regions.

The derived polynucleotide will not necessarily be derived physically from the nucleotide sequence of interest  
35 under study, but may be generated in any manner, including but not limited to chemical synthesis, replication, reverse transcription or transcription, which is based on the

information provided by the sequence of bases in the region(s) from which the polynucleotide is derived; as such, it may represent either a sense or an antisense orientation of the original polynucleotide. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with an intended use.

The term "primer" denotes a specific oligonucleotide sequence complementary to a target nucleotide sequence and used to hybridize to the target nucleotide sequence and to serve as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase.

The term "probe" denotes a defined nucleic acid segment (or nucleotide analog segment, i.e., PNA) which can be used to identify specific DNA present in samples bearing the complementary sequence.

A "polypeptide" or "amino acid" sequence derived from a designated nucleic acid sequence refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence or a portion thereof wherein the portion consists of at least 3 to 5 amino acids, and more preferably at least 8 to 10 amino acids, and even more preferably 15 to 20 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A "recombinant polypeptide" as used herein means at least a polypeptide which, by virtue of its origin or manipulation, is not associated with all or a portion of the polypeptide with which it is associated in nature and/or is linked to a polypeptide other than that to which it is linked in nature. A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence. It also may be generated in any manner, including chemical synthesis or expression by a recombinant expression system.

The term "synthetic peptide" as used herein means a polymeric form of amino acids of any length, which may be chemically synthesized by methods well-known to the routineer. These synthetic peptides are useful in various applications.

The term "polynucleotide" as used herein means a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, the term includes double- and single-stranded DNA, as well as double- and single-stranded RNA. It also includes modifications, such as methylation or capping, and unmodified forms of the polynucleotide.

"A sequence corresponding to a cDNA" means that the sequence contains a polynucleotide sequence that is identical to or complementary to a sequence in the designated DNA. The degree (or "percent") of identity or complementarity to the cDNA will be approximately 50% or greater, will preferably be at least about 70% or greater, and more preferably will be at least about 90%. The sequence that corresponds will be at least about 50 nucleotides in length, will preferably be about 60 nucleotides in length, and more preferably, will be at least about 70 nucleotides in length. The correspondence between the gene or gene fragment of interest and the cDNA can be determined by methods known in the art, which include, for example, a direct comparison of the sequenced material with the cDNAs described, or hybridization and digestion with single strand nucleases, followed by size determination of the digested fragments.

"Purified polynucleotide" refers to a polynucleotide of interest or fragment thereof which is essentially free, i.e., contains less than about 50%, preferably less than about 70%, and more preferably, less than about 90% of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example,

disruption of the cell containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

5 Thus, "purified polypeptide" means a polypeptide of interest or fragment thereof which is essentially free, that is, contains less than about 50%, preferably less than about 70%, and more preferably, less than about 90% of cellular components with which the polypeptide of interest is  
10 naturally associated. Methods for purifying are known in the art.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-  
15 occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, which is separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such  
20 polynucleotide or polypeptide could be part of a composition, and still be isolated in that the vector or composition is not part of its natural environment.

"Polypeptide" as used herein indicates a molecular chain of amino acids and does not refer to a specific length  
25 of the product. Thus, peptides, oligopeptides and proteins are included within the definition of polypeptide. This term, however, is not intended to refer to post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like.

30 "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vectors or other  
35 transferred DNA, and include the original progeny of the original cell which has been transfected.

As used herein "replicon" means any genetic element, such as a plasmid, a chromosome or a virus, that behaves as an autonomous unit of polynucleotide replication within a cell.

5       A "vector" is a replicon in which another polynucleotide segment is attached, such as to bring about the replication and/or expression of the attached segment.

10       The term "control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, such control sequences generally include promoter, ribosomal binding site and terminators; in eukaryotes, such control sequences generally include  
15       promoters, terminators and, in some instances, enhancers. The term "control sequence" thus is intended to include at a minimum all components whose presence is necessary for expression, and also may include additional components whose presence is advantageous, for example, leader sequences.

20       "Operably linked" refers to a situation wherein the components described are in a relationship permitting them to function in their intended manner. Thus, for example, a control sequence "operably linked" to a coding sequence is ligated in such a manner that expression of the coding  
25       sequence is achieved under conditions compatible with the control sequences.

30       The term "open reading frame" or "ORF" refers to a region of a polynucleotide sequence which encodes a polypeptide; this region may represent a portion of a coding sequence or a total coding sequence.

35       A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5' -terminus and a translation stop codon at the 3' -terminus. A coding

sequence can include, but is not limited to, mRNA, cDNA, and recombinant polynucleotide sequences.

The term "immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptide(s) which also  
5 are present in and are unique to the designated polypeptide(s). Immunological identity may be determined by antibody binding and/or competition in binding. These techniques are known to the routineer and also are described herein. The uniqueness of an epitope also can be determined  
10 by computer searches of known data banks, such as GenBank, for the polynucleotide sequences which encode the epitope, and by amino acid sequence comparisons with other known proteins.

As used herein, "epitope" means an antigenic  
15 determinant of a polypeptide. Conceivably, an epitope can comprise three amino acids in a spatial conformation which is unique to the epitope. Generally, an epitope consists of at least five such amino acids, and more usually, it consists of at least eight to ten amino acids. Methods of  
20 examining spatial conformation are known in the art and include, for example, x-ray crystallography and two-dimensional nuclear magnetic resonance.

A "conformational epitope" is an epitope that is comprised of a specific juxtaposition of amino acids in an  
25 immunologically recognizable structure, such amino acids being present on the same polypeptide in a contiguous or non-contiguous order or present on different polypeptides.

A polypeptide is "immunologically reactive" with an antibody when it binds to an antibody due to antibody  
30 recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly by the kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an epitope  
35 against which the antibody is directed. The methods for determining whether a polypeptide is immunologically reactive with an antibody are known in the art.

As used herein, the term "immunogenic polypeptide containing an epitope of interest" means naturally occurring polypeptides of interest or fragments thereof, as well as polypeptides prepared by other means, for example, by  
5 chemical synthesis or the expression of the polypeptide in a recombinant organism.

The term "transformation" refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion. For example, direct  
10 uptake, transduction or f-mating are included. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

"Treatment" refers to prophylaxis and/or therapy.

15 The term "individual" as used herein refers to vertebrates, particularly members of the mammalian species and includes but is not limited to domestic animals, sports animals, primates and humans; more particularly the term refers to humans.

20 The term "sense strand" or "plus strand" (or "+") as used herein denotes a nucleic acid that contains the sequence that encodes the polypeptide. The term "antisense strand" or "minus strand" (or "-") denotes a nucleic acid that contains a sequence that is complementary to that of  
25 the "plus" strand.

The term "test sample" refers to a component of an individual's body which is the source of the analyte (such as, antibodies of interest or antigens of interest). These components are well known in the art. These test samples  
30 include biological samples which can be tested by the methods of the present invention described herein and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, urine, lymph fluids, and various external secretions of the respiratory, intestinal  
35 and genitorurinary tracts, tears, saliva, milk, white blood cells, myelomas and the like; biological fluids such as cell



culture supernatants; fixed tissue specimens; and fixed cell specimens.

"Purified product" refers to a preparation of the product which has been isolated from the cellular  
5 constituents with which the product is normally associated, and from other types of cells which may be present in the sample of interest.

"PNA" denotes a "peptide nucleic acid analog" which may be utilized in a procedure such as an assay described herein  
10 to determine the presence of a target. "MA" denotes a "morpholino analog" which may be utilized in a procedure such as an assay described herein to determine the presence of a target. See, for example, U.S. Patent No. 5,378,841, which is incorporated herein by reference. PNAs are  
15 neutrally charged moieties which can be directed against RNA targets or DNA. PNA probes used in assays in place of, for example, the DNA probes of the present invention, offer advantages not achievable when DNA probes are used. These advantages include manufacturability, large scale labeling, reproducibility, stability, insensitivity to changes in  
20 ionic strength and resistance to enzymatic degradation which is present in methods utilizing DNA or RNA. These PNAs can be labeled with such signal generating compounds as fluorescein, radionucleotides, chemiluminescent compounds, and the like. PNAs or other nucleic acid analogs such as  
25 MAs thus can be used in assay methods in place of DNA or RNA. Although assays are described herein utilizing DNA probes, it is within the scope of the routineer that PNAs or MAs can be substituted for RNA or DNA with appropriate  
30 changes if and as needed in assay reagents.

"Analyte," as used herein, is the substance to be detected which may be present in the test sample. The analyte can be any substance for which there exists a naturally occurring specific binding member (such as, an  
35 antibody), or for which a specific binding member can be prepared. Thus, an analyte is a substance that can bind to one or more specific binding members in an assay. "Analyte"

also includes any antigenic substances, haptens, antibodies, and combinations thereof. As a member of a specific binding pair, the analyte can be detected by means of naturally occurring specific binding partners (pairs) such as the use of intrinsic factor protein as a member of a specific binding pair for the determination of Vitamin B12, the use of folate-binding protein to determine folic acid, or the use of a lectin as a member of a specific binding pair for the determination of a carbohydrate. The analyte can include a protein, a peptide, an amino acid, a nucleotide target, and the like.

"Inflammation" or "inflammatory disease," as used herein, refer to infiltration of activated lymphocytes such as neutrophils, eosinophils, macrophages, T cells and B-cells, into a host tissue that results in damage to the host organism. Examples of inflammatory disease include but are not limited to conditions such as inflammatory bowel disease, sepsis, and rheumatoid arthritis.

An "Expressed Sequence Tag" or "EST" refers to the partial sequence of a cDNA insert which has been made by reverse transcription of mRNA extracted from a tissue, followed by insertion into a vector.

A "transcript image" refers to a table or list giving the quantitative distribution of ESTs in a library and represents the genes active in the tissue from which the library was made.

The present invention provides assays which utilize specific binding members. A "specific binding member," as used herein, is a member of a specific binding pair. That is, two different molecules where one of the molecules through chemical or physical means specifically binds to the second molecule. Therefore, in addition to antigen and antibody specific binding pairs of common immunoassays, other specific binding pairs can include biotin and avidin, carbohydrates and lectins, complementary nucleotide sequences, effector and receptor molecules, cofactors and enzymes, enzyme inhibitors and enzymes, and the like.

Furthermore, specific binding pairs can include members that are analogs of the original specific binding members, for example, an analyte-analog. Immunoreactive specific binding members include antigens, antigen fragments, antibodies and  
5 antibody fragments, both monoclonal and polyclonal, and complexes thereof, including those formed by recombinant DNA molecules.

The term "hapten," as used herein, refers to a partial antigen or non-protein binding member which is capable of  
10 binding to an antibody, but which is not capable of eliciting antibody formation unless coupled to a carrier protein.

A "capture reagent," as used herein, refers to an unlabeled specific binding member which is specific either  
15 for the analyte as in a sandwich assay, for the indicator reagent or analyte as in a competitive assay, or for an ancillary specific binding member, which itself is specific for the analyte, as in an indirect assay. The capture reagent can be directly or indirectly bound to a solid phase  
20 material before the performance of the assay or during the performance of the assay, thereby enabling the separation of immobilized complexes from the test sample.

The "indicator reagent" comprises a "signal-generating compound" ("label") which is capable of generating and  
25 generates a measurable signal detectable by external means, conjugated ("attached") to a specific binding member.

"Specific binding member" as used herein means a member of a specific binding pair. That is, two different molecules where one of the molecules through chemical or physical  
30 means specifically binds to the second molecule. In addition to being an antibody member of a specific binding pair, the indicator reagent also can be a member of any specific binding pair, including either hapten-anti-hapten systems such as biotin or anti-biotin, avidin or biotin, a  
35 carbohydrate or a lectin, a complementary nucleotide sequence, an effector or a receptor molecule, an enzyme cofactor and an enzyme, an enzyme inhibitor or an enzyme,

and the like. An immunoreactive specific binding member can be an antibody, an antigen, or an antibody/antigen complex that is capable of binding either to a polypeptide of interest as in a sandwich assay, to the capture reagent as  
5 in a competitive assay, or to the ancillary specific binding member as in an indirect assay. When describing probes and probe assays, the term "reporter molecule" may be used. A reporter molecule comprises a signal generating compound as described hereinabove conjugated to a specific binding  
10 member of a specific binding pair, such as carbazol or adamantane.

The various "signal-generating compounds" (labels) contemplated include chromogens, catalysts such as enzymes, luminescent compounds such as fluorescein and rhodamine,  
15 chemiluminescent compounds such as dioxetanes, acridiniums, phenanthridiniums and luminol, radioactive elements, and direct visual labels. Examples of enzymes include alkaline phosphatase, horseradish peroxidase, beta-galactosidase, and the like. The selection of a particular label is not  
20 critical, but it will be capable of producing a signal either by itself or in conjunction with one or more additional substances.

"Solid phases" ("solid supports") are known to those in the art and include the walls of wells of a reaction tray,  
25 test tubes, polystyrene beads, magnetic beads, nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, and Duracytes® (red blood cells "fixed" by pyruvic aldehyde and formaldehyde, available from Abbott Laboratories, Abbott  
30 Park, IL) and others. The "solid phase" is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red  
35 blood cells and Duracytes® are all suitable examples. Suitable methods for immobilizing peptides on solid phases include ionic, hydrophobic, covalent interactions and the

like. A "solid phase", as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid phase can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid phase and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid phase material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, Duracytes® and other configurations known to those of ordinary skill in the art.

It is contemplated and within the scope of the present invention that the solid phase also can comprise any suitable porous material with sufficient porosity to allow access by detection antibodies and a suitable surface affinity to bind antigens. Microporous structure generally are preferred, but materials with gel structure in the hydrated state may be used as well. Such useful solid supports include but are not limited to nitrocellulose and nylon. It is contemplated that such porous solid supports described herein preferably are in the form of sheets of thickness from about 0.01 to 0.5 mm, preferably about 0.1mm. The pore size may vary within wide limits, and preferably is from about 0.025 to 15 microns, especially from about 0.15 to 15 microns. The surface of such supports may be

activated by chemical processes which cause covalent linkage of the antigen or antibody to the support. The irreversible binding of the antigen or antibody is obtained, however, in general, by adsorption on the porous material by poorly understood hydrophobic forces. Other suitable solid supports are known in the art.

#### Reagents

The present invention provides reagents such as polynucleotide sequences derived from a TREPA of interest, polypeptides encoded thereby, and antibodies developed from these polypeptides. The present invention also provides reagents such as oligonucleotide fragments derived from the disclosed polynucleotides and nucleic acid sequences complementary to the these polynucleotides. The polynucleotides or polypeptides or antibodies of the present invention may be used in the diagnosis, prognosis, and/or treatment of individuals with conditions associated with the TREPA gene, such as inflammation, or to identify a predisposition to this condition. The sequences disclosed herein represent unique polynucleotides which can be used in assays or for producing a specific profile of gene transcription activity.

Selected TREPA-derived polynucleotides can be used in the methods described herein for the detection of normal or altered gene expression. Such methods may employ the TREPA-derived polynucleotides disclosed herein or oligonucleotides, fragments or derivatives thereof, or nucleic acid sequences complementary to these polynucleotides.

The polynucleotides disclosed herein, their complementary sequences or fragments of either can be used in assays to detect, amplify or quantify genes, cDNAs or mRNAs relating to TREPA disease and conditions associated with it. They also can be used to identify an entire or partial coding region which encodes a TREPA polypeptide. They further can be provided in individual containers in the form of a kit for assays, or provided as individual

compositions. If provided in a kit for assays, other suitable reagents such as buffers, conjugates and the like may be included.

The polynucleotide(s) may be in the form of mRNA or  
5 DNA. Polynucleotides in the form of DNA, cDNA, genomic DNA, and synthetic DNA are within the scope of the present invention. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding (sense) strand or non-coding (anti-sense) strand. The coding  
10 sequence which encodes the polypeptide may be identical to the coding sequence provided herein or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same polypeptide as the DNA provided herein.

15 This polynucleotide may include only the coding sequence for the polypeptide, or the coding sequence for the polypeptide and additional coding sequence such as a leader or secretory sequence or a proprotein sequence, or the coding sequence for the polypeptide (and optionally  
20 additional coding sequence) and non-coding sequence, such as a non-coding sequence 5' and/or 3' of the coding sequence for the polypeptide.

In addition, the invention includes variant polynucleotides containing modifications such as  
25 polynucleotide deletions, substitutions or additions; and any polypeptide modification resulting from the variant polynucleotide sequence. A polynucleotide of the present invention also may have a coding sequence which is a naturally occurring allelic variant of the coding sequence  
30 provided herein.

In addition, the coding sequence for the polypeptide may be fused in the same reading frame to a polynucleotide sequence which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence  
35 which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the

leader sequence cleaved by the host cell to form the polypeptide. The polynucleotides may also encode for a proprotein which is the protein plus additional 5' amino acid residues. A protein having a prosequence is a  
5 proprotein and may in some cases be an inactive form of the protein. Once the prosequence is cleaved, an active protein remains. Thus, the polynucleotide of the present invention may encode for a protein, or for a protein having a prosequence or for a protein having both a presequence  
10 (leader sequence) and a prosequence.

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexa-  
15 histidine tag supplied by a pQE-9 vector to provide for purification of the polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g. COS-7 cells, is used. The HA tag corresponds to  
20 an epitope derived from the influenza hemagglutinin protein. See, for example, I. Wilson, et al., Cell 37:767 (1984).

It is contemplated that polynucleotides will be considered to hybridize to the sequences provided herein if there is at least 50%, and preferably at least 70%, identity  
25 between the polynucleotide and the sequence.

The present invention also provides an antibody produced by using a purified TREPA gene polypeptide of which at least a portion of the polypeptide is encoded by the TREPA gene polynucleotide selected from the polynucleotides  
30 provided herein. These antibodies may be used in the methods provided herein for the detection of TREPA polypeptides in test samples. The antibody also may be used for therapeutic purposes, for example, in neutralizing the activity of a TREPA polypeptide in conditions associated  
35 with altered or abnormal expression.

The present invention further relates to a TREPA polypeptide which has the deduced amino acid sequence as



provided herein, as well as fragments, analogs and derivatives of such a polypeptide. The polypeptide of the present invention may be a recombinant polypeptide, a natural purified polypeptide or a synthetic polypeptide.

5 The fragment, derivative or analog of the TREPA polypeptide may be one in which one or more of the amino acid residues is substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded  
10 by the genetic code; or it may be one in which one or more of the amino acid residues includes a substituent group; or it may be one in which the polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol); or it  
15 may be one in which the additional amino acids are fused to the polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are within the scope of the present  
20 invention. The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably purified.

Thus, a polypeptide of the present invention may have an amino acid sequence that is identical to that of the  
25 naturally occurring polypeptide or that is different by minor variations due to one or more amino acid substitutions. The variation may be a "conservative change" typically in the range of about 1 to 5 amino acids, wherein the substituted amino acid has similar structural or  
30 chemical properties, e.g., replacement of leucine with isoleucine or threonine with serine. In contrast, variations may include nonconservative changes, e.g., replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or  
35 insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without changing biological or immunological

activity may be found using computer programs well known in the art, for example, DNASTAR software (DNASTAR Inc., Madison WI).

Probes constructed according to the polynucleotide sequences of the present invention can be used in various assay methods to provide various types of analysis. For example, such probes can be used in Fluorescent In Situ Hybridization (FISH) technology to perform chromosomal analysis, and used to identify cancer-specific structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR-generated and/or allele specific oligonucleotides probes, allele specific amplification or by direct sequencing. Probes also can be labeled with radioisotopes, directly- or indirectly- detectable haptens, or fluorescent molecules, and utilized for in situ hybridization studies to evaluate the mRNA expression of the gene comprising the polynucleotide in fixed tissue specimens or cells.

This invention also provides teachings as to the production of the polynucleotides and polypeptides provided herein.

#### Probe Assays

The sequences provided herein may be used to produce probes which can be used in assays for the detection of nucleic acids in test samples. The probes may be designed from conserved nucleotide regions of the polynucleotides of interest or from non-conserved nucleotide regions of the polynucleotides of interest. The design of such probes for optimization in assays is within the skill of the routineer. Generally, nucleic acid probes are developed from non-conserved or unique regions when maximum specificity is desired, and nucleic acid probes are developed from conserved regions when assaying for nucleotide regions that are closely related to, for example, different members of a multigene family or in related species like mouse and man.

The polymerase chain reaction (PCR) is a technique for amplifying a desired nucleic acid sequence (target) contained in a nucleic acid or mixture thereof. In PCR, a pair of primers is employed in excess to hybridize at the outside ends of complementary strands of the target nucleic acid. The primers are each extended by a polymerase using the target nucleic acid as a template. The extension products become target sequences themselves, following dissociation from the original target strand. New primers then are hybridized and extended by a polymerase, and the cycle is repeated to geometrically increase the number of target sequence molecules. PCR is disclosed in U.S. patents 4,683,195 and 4,683,202, which are incorporated herein by reference.

The Ligase Chain Reaction (LCR) is an alternate method for nucleic acid amplification. In LCR, probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of

hybridization and ligation, amplification of the target sequence is achieved. This technique is described more completely in EP-A- 320 308 to K. Backman published June 16, 1989 and EP-A-439 182 to K. Backman et al., published July 31, 1991, both of which are incorporated herein by reference.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Patent No. 5,322,770, which is incorporated herein by reference; or reverse transcribe mRNA into cDNA followed by asymmetric gap ligase chain reaction (RT-AGLCR) as described by R. L. Marshall, et al., PCR Methods and Applications 4: 80-84 (1994), which also is incorporated herein by reference.

Other known amplification methods which can be utilized herein include but are not limited to the so-called "NASBA" or "3SR" technique described in PNAS USA 87:1874-1878 (1990) and also described in Nature 350 (No. 6313):91-92 (1991); Q-beta amplification as described in published European Patent Application (EPA) No. 4544610; strand displacement amplification (as described in G. T. Walker et al., Clin. Chem. 42:9-13 (1996) and European Patent Application No. 684315; and target mediated amplification, as described by PCT Publication WO 9322461.

In one embodiment, the present invention generally comprises the steps of contacting a test sample suspected of containing a target polynucleotide sequence with amplification reaction reagents comprising an amplification primer, and a detection probe that can hybridize with an internal region of the amplicon sequences. Probes and primers employed according to the method herein provided are labeled with capture and detection labels wherein probes are labeled with one type of label and primers are labeled with the other type of label. Additionally, the primers and probes are selected such that the probe sequence has a lower melt temperature than the primer sequences. The

amplification reagents, detection reagents and test sample are placed under amplification conditions whereby, in the presence of target sequence, copies of the target sequence (an amplicon) are produced. In the usual case, the amplicon is double stranded because primers are provided to amplify a target sequence and its complementary strand. The double stranded amplicon then is thermally denatured to produce single stranded amplicon members. Upon formation of the single stranded amplicon members, the mixture is cooled to allow the formation of complexes between the probes and single stranded amplicon members.

After the probe/single stranded amplicon member hybrids are formed, they are detected. Standard heterogeneous assay formats are suitable for detecting the hybrids using the detection labels and capture labels present on the primers and probes. The hybrids can be bound to a solid phase reagent by virtue of the capture label and detected by virtue of the detection label. In cases where the detection label is directly detectable, the presence of the hybrids on the solid phase can be detected by causing the label to produce a detectable signal, if necessary, and detecting the signal. In cases where the label is not directly detectable, the captured hybrids can be contacted with a conjugate, which generally comprises a binding member attached to a directly detectable label. The conjugate becomes bound to the complexes, and the conjugate's presence on the complexes can be detected with the directly detectable label. Thus, the presence of the hybrids on the solid phase reagent can be determined. Those skilled in the art will recognize that wash steps may be employed to wash away unhybridized amplicon or probe as well as unbound conjugate.

A test sample is typically anything suspected of containing a target sequence. Test samples can be prepared using methodologies well known in the art such as by obtaining a specimen from an individual and, if necessary, disrupting any cells contained therein to release target

nucleic acids. Although the target sequence is described as single stranded, it also is contemplated to include the case where the target sequence is actually double stranded but is merely separated from its complement prior to hybridization with the amplification primer sequences. In the case where PCR is employed in this method, the ends of the target sequences are usually known. In cases where LCR or a modification thereof is employed in the preferred method, the entire target sequence is usually known. Typically, the target sequence is a nucleic acid sequence such as, for example, RNA or DNA.

The method provided herein can be used in well known amplification reactions that include thermal cycle reaction mixtures, particularly in PCR and GLCR. Amplification reactions typically employ primers to repeatedly generate copies of a target nucleic acid sequence, which target sequence is usually a small region of a much larger nucleic acid sequence. Primers are themselves nucleic acid sequences that are complementary to regions of a target sequence. Under amplification conditions, these primers hybridize or bind to the complementary regions of the target sequence. Copies of the target sequence typically are generated by the process of primer extension and/or ligation which utilizes enzymes with polymerase or ligase activity, separately or in combination, to add nucleotides to the hybridized primers and/or ligate adjacent probe pairs. The nucleotides that are added to the primers or probes, as monomers or preformed oligomers, are also complementary to the target sequence. Once the primers or probes have been sufficiently extended and/or ligated, they are separated from the target sequence, for example, by heating the reaction mixture to a "melt temperature" which is one in which complementary nucleic acid strands dissociate. Thus, a sequence complementary to the target sequence is formed.

A new amplification cycle then can take place to further amplify the number of target sequences by separating any double stranded sequences, allowing primers or probes to

hybridize to their respective targets, extending and/or ligating the hybridized primers or probes and re-separating. The complementary sequences that are generated by amplification cycles can serve as templates for primer  
5 extension or filling the gap of two probes to further amplify the number of target sequences. Typically, a reaction mixture is cycled between 20 and 100 times, more typically, a reaction mixture is cycled between 25 and 50 times. The numbers of cycles can be determined by the  
10 routineer. In this manner, multiple copies of the target sequence and its complementary sequence are produced. Thus, primers initiate amplification of the target sequence when it is present under amplification conditions.

Generally, two primers which are complementary to a  
15 portion of a target strand and its complement are employed in PCR. For LCR, four probes, two of which are complementary to a target sequence and two of which are similarly complementary to the target's complement, are generally employed. In addition to the primer sets and  
20 enzymes previously mentioned, a nucleic acid amplification reaction mixture may also comprise other reagents which are well known and include but are not limited to: enzyme cofactors such as manganese; magnesium; salts; nicotinamide adenine dinucleotide (NAD); and deoxynucleotide  
25 triphosphates (dNTPs) such as for example deoxyadenine triphosphate, deoxyguanine triphosphate, deoxycytosine triphosphate and deoxythymine triphosphate.

While the amplification primers initiate amplification of the target sequence, the detection (or hybridization)  
30 probe is not involved in amplification. Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example, peptide nucleic acids which are disclosed in International Patent Application WO 92/20702; morpholino analogs which are described in U.S.  
35 Patents Nos. 5,185,444, 5,034,506, and 5,142,047; and the like. Depending upon the type of label carried by the probe, the probe is employed to capture or detect the

amplicon generated by the amplification reaction. The probe is not involved in amplification of the target sequence and therefore may have to be rendered "non-extendible" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendible and nucleic acid probes can be rendered non-extendible by modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified. U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 and incorporated herein by reference describes modifications which can be used to render a probe non-extendible.

Accordingly, the ratio of primers to probes is not important. Thus, either the probes or primers can be added to the reaction mixture in excess whereby the concentration of one would be greater than the concentration of the other. Alternatively, primers and probes can be employed in equivalent concentrations. Preferably, however, the primers are added to the reaction mixture in excess of the probes. Thus, primer to probe ratios of, for example, 5:1 and 20:1 are preferred.

While the length of the primers and probes can vary, the probe sequences are selected such that they have a lower melt temperature than the primer sequences. Hence, the primer sequences are generally longer than the probe sequences. Typically, the primer sequences are in the range of between 20 and 50 nucleotides long, more typically in the range of between 20 and 30 nucleotides long. The typical probe is in the range of between 10 and 25 nucleotides long.

Various methods for synthesizing primers and probes are well known in the art. Similarly, methods for attaching labels to primers or probes are also well known in the art. For example, it is a matter of routine to synthesize desired



nucleic acid primers or probes using conventional nucleotide phosphoramidite chemistry and instruments available from Applied Biosystems, Inc., (Foster City, CA), Dupont (Wilmington, DE), or Milligen (Bedford MA). Many methods  
5 have been described for labeling oligonucleotides such as the primers or probes of the present invention. Enzo Biochemical (New York, NY) and Clontech (Palo Alto, CA) both have described and commercialized probe labeling techniques. For example, a primary amine can be attached to a 3' oligo  
10 terminus using 3'-Amine-ON CPG™ (Clontech, Palo Alto, CA). Similarly, a primary amine can be attached to a 5' oligo terminus using Aminomodifier II® (Clontech). The amines can be reacted to various haptens using conventional activation and linking chemistries. In addition, copending  
15 applications U.S. Serial Nos. 625,566, filed December 11, 1990 and 630,908, filed December 20, 1990, which are each incorporated herein by reference, teach methods for labeling probes at their 5' and 3' termini, respectively. Publications WO92/10505, published 25 June 1992 and WO  
20 92/11388 published 9 July 1992 teach methods for labeling probes at their 5' and 3' ends, respectively. According to one known method for labeling an oligonucleotide, a label-phosphoramidite reagent is prepared and used to add the label to the oligonucleotide during its synthesis. See, for  
25 example, N.T. Thuong et al., Tet. Letters 29(46):5905-5908 (1988); or J. S. Cohen et al., published U.S. Patent Application 07/246,688 (NTIS ORDER No. PAT-APPL-7-246,688) (1989). Preferably, probes are labeled at their 3' and 5' ends.

30 Capture labels are carried by the primers or probes and can be a specific binding member which forms a binding pair with the solid phase reagent's specific binding member. It will be understood, of course, that the primer or probe itself may serve as the capture label. For example, in the  
35 case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of the primer or probe to thereby

immobilize the primer or probe to the solid phase. In cases where the probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the single stranded amplicon members. In the case where the primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase because the probe is selected such that it is not fully complementary to the primer sequence.

Generally, probe/single stranded amplicon member complexes can be detected using techniques commonly employed to perform heterogeneous immunoassays. Preferably, in this embodiment, detection is performed according to the protocols used by the commercially available Abbott LCx<sup>®</sup> instrumentation (Abbott Laboratories, Abbott Park, IL).

The primers and probes disclosed herein are useful in typical PCR assays, wherein the test sample is contacted with a pair of primers, amplification is performed, the hybridization probe is added, and detection is performed.

Another method provided by the present invention comprises contacting a test sample with a plurality of polynucleotides wherein at least one polynucleotide is provided herein, hybridizing the test sample with the plurality of polynucleotides and detecting the hybridization complexes. The hybridization complexes are identified and quantitated to compile a profile which is indicative of TREPA disease. Expressed RNA sequences may further be detected by reverse transcription and amplification of the DNA product by procedures well-known in the art, including the polymerase chain reaction (PCR).

#### Drug Screening and Gene Therapy

The present invention also encompasses the use of gene therapy methods for the introduction of anti-sense TREPA gene-derived molecules such as polynucleotides or oligonucleotides of the present invention into patients with conditions associated with abnormal expression of

polynucleotides related to TREPA disease including cancer. These molecules, including antisense RNA and DNA fragments and ribozymes, are designed to inhibit the translation of a TREPA-derived polynucleotide mRNA, and may be used  
5 therapeutically in the treatment of conditions associated with altered or abnormal expression of a TREPA-derived polynucleotide.

Alternatively, the oligonucleotides described above can be delivered to cells by procedures in the art such that the  
10 anti-sense RNA or DNA may be expressed in vivo to inhibit production of TREPA-derived polypeptide in the manner described above. Antisense constructs to TREPA-derived polynucleotide, therefore, reverse the action of TREPA-derived transcripts and may be used for treating TREPA  
15 disease conditions, such as inflammation. These antisense constructs may also be used to treat tumor metastases.

Effects on tumor vasculature, such as those associated with this family of molecules (M. W. Boehme, Eur. J. Clin. Invest. 26: 404-410 1996), are useful in the treatment of  
20 both primary and metastatic solid tumors, including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder and urothelium), female genital  
25 tract, (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate, seminal vesicles, testes and and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well  
30 as hemangiomas, melanomas, sarcomas (including those arising from bone and soft tissues as well as Kaposi's sarcoma) and tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas).  
35 Such proteins may also be useful in treating solid tumors arising from hematopoietic malignancies such as leukemias (i.e., chloromas, plasmacytomas and the plaques and tumors

of mycosis fungoides and cutaneous T-cell lymphoma/leukemia) as well as in the treatment of lymphomas (both Hodgkin's and non-Hodgkin's lymphomas). In addition, these proteins or genes which encode their expression may be useful in the prevention of metastases from the tumors described above either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

Further uses include the treatment and prophylaxis of autoimmune diseases such as rheumatoid, immune and degenerative arthritis; various ocular diseases such as diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, retrolental fibroplasia, neovascular glaucoma, rubeosis, retinal neovascularization due to macular degeneration, hypoxia, and other abnormal neovascularization conditions of the eye; skin diseases such as psoriasis; blood vessel diseases such as hemangiomas, and capillary proliferation within atherosclerotic plaques; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation. Other uses include the treatment of diseases characterized by excessive or abnormal stimulation of endothelial cells, including but not limited to intestinal adhesions, Crohn's disease, atherosclerosis, scleroderma, and hypertrophic scars, i.e. keloids. Another use is as a birth control agent, by inhibiting ovulation and establishment of the placenta. TREPA is also useful in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (Rochele minalia quintosa) and ulcers (Helicobacter pylori).

TREPA may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with TREPA and then TREPA may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor. Additionally, TREPA, TREPA fragments, TREPA antisera, TREPA

receptor agonists, TREPA receptor antagonists, or combinations thereof, may be combined with pharmaceutically acceptable excipients, and optionally sustained-release matrices, such as biodegradable polymers, to form

5 therapeutic compositions.

A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid-base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes  
10 and body fluids. A sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides,  
15 poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene,  
20 polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (copolymers of lactic acid and glycolic acid).

Cytotoxic agents such as ricin, may be linked to TREPA,  
25 and high affinity TREPA peptide fragments, thereby providing a tool for destruction of cells that bind TREPA. Peptides linked to cytotoxic agents may be infused in a manner designed to maximize delivery to the desired location. For example, ricin-linked high affinity TREPA fragments may be  
30 delivered through a cannula into vessels supplying the target site or directly into the target. Such agents may also be delivered in a controlled manner through osmotic pumps coupled to infusion cannulae. A combination of TREPA antagonists may be co-applied with stimulators of  
35 angiogenesis to increase vascularization of tissue. Therapeutic regimens of this type could provide an effective means of destroying metastatic cancer.

The present invention also encompasses gene therapy whereby the gene encoding TREPA is regulated in a patient. Various methods of transferring or delivering DNA to cells for expression of the gene product protein, otherwise  
5 referred to as gene therapy, are disclosed in "Gene Transfer into Mammalian Somatic Cells in vivo", N. Yang, Crit. Rev. Biotechn. 12(4): 335-356 (1992), which is hereby incorporated by reference. Gene therapy encompasses incorporation of DNA sequences into somatic cells or germ  
10 line cells for use in either ex vivo or in vivo therapy. Gene therapy functions to replace genes, augment normal or abnormal gene function, and to combat infectious diseases and other pathologies.

Strategies for treating these medical problems with  
15 gene therapy include therapeutic strategies such as identifying the defective gene and then adding a functional gene to either replace the function of the defective gene or to augment a slightly functional gene; or prophylactic strategies, such as adding a gene which encodes a protein  
20 product that will treat the condition or that will make the tissue or organ more susceptible to a treatment regimen. As an example of a prophylactic strategy, a gene encoding TREPA may be placed in a patient and thus prevent occurrence of angiogenesis; or a gene that makes tumor cells more  
25 susceptible to radiation could be inserted so that radiation of the tumor would cause increased killing of the tumor cells.

Many protocols for transfer of TREPA DNA or TREPA regulatory sequences are envisioned in this invention.  
30 Transfection of promoter sequences, other than one specifically associated with TREPA, or other sequences which would increase production of TREPA protein are also envisioned as methods of gene therapy. An example of this technology is found in Transkaryotic Therapies, Inc., of  
35 Cambridge, Massachusetts, using homologous recombination to insert a "genetic switch" that turns on an erythropoietin gene in cells. See Genetic Engineering News, April 15,

0902700 004000

1994. Such "genetic switches" could be used to activate TREPA (or a TREPA receptor) in cells not normally expressing these proteins.

Gene transfer methods for gene therapy fall into three  
5 broad categories: (1) physical (e.g., electroporation,  
direct gene transfer and particle bombardment), (2)  
chemical (e.g., lipid-based carriers and other non-viral  
vectors) and (3) biological (e.g., virus derived vectors).  
For example, non-viral vectors such as liposomes coated with  
10 DNA may be directly injected intravenously into the patient.  
It is believed that the liposome/DNA complexes are  
concentrated in the liver where they deliver the DNA to  
macrophages and Kupffer cells. Additionally, vectors or the  
"naked" DNA of the gene may be directly injected into the  
15 desired organ, tissue or tumor for targeted delivery of the  
therapeutic DNA.

Gene therapy methodologies can also be described by  
delivery site. Fundamental ways to deliver genes include ex  
vivo gene transfer, in vivo gene transfer, and in vitro gene  
20 transfer. In ex vivo gene transfer, cells are taken from  
the patient and grown in cell culture. The DNA is  
transfected into the cells, the transfected cells are  
expanded in number and then reimplanted in the patient. In  
in vitro gene transfer, the transformed cells are cells  
25 growing in culture, such as tissue culture cells, and not  
particular cells from a particular patient. These  
"laboratory cells" are transfected, and the transfected  
cells are selected and expanded for either implantation into  
a patient or for other uses. In vivo gene transfer involves  
30 introducing the DNA into the cells of the patient when the  
cells are within the patient. All three of the broad based  
categories described above may be used to achieve gene  
transfer in vivo, ex vivo, and in vitro.

Mechanical (i.e. physical) methods of DNA delivery can  
35 be achieved by direct injection of DNA, such as  
microinjection of DNA into germ or somatic cells,  
pneumatically delivered DNA-coated particles, such as the

gold particles used in a "gene gun," and inorganic chemical approaches such as calcium phosphate transfection. It has been found that physical injection of plasmid DNA into muscle cells yields a high percentage of cells which are  
5 transfected and have a sustained expression of marker genes. The plasmid DNA may or may not integrate into the genome of the cells. Non-integration of the transfected DNA would allow the transfection and expression of gene product proteins in terminally differentiated, non-proliferative  
10 tissues for a prolonged period of time without fear of mutational insertions, deletions, or alterations in the cellular or mitochondrial genome. Long-term, but not necessarily permanent, transfer of therapeutic genes into specific cells may provide treatments for genetic diseases  
15 or for prophylactic use. The DNA could be reinjected periodically to maintain the gene product level without mutations occurring in the genomes of the recipient cells. Non-integration of exogenous DNAs may allow for the presence of several different exogenous DNA constructs within one  
20 cell with all of the constructs expressing various gene products.

Particle-mediated gene transfer may also be employed for injecting DNA into cells, tissues and organs. With a particle bombardment device, or "gene gun," a motive force  
25 is generated to accelerate DNA-coated high density particles (such as gold or tungsten) to a high velocity that allows penetration of the target organs, tissues or cells. Electroporation for gene transfer uses an electrical current to make cells or tissues susceptible to electroporation-  
30 mediated gene transfer. A brief electric impulse with a given field strength is used to increase the permeability of a membrane in such a way that DNA molecules can penetrate into the cells. The techniques of particle-mediated gene transfer and electroporation are well known to those of  
35 ordinary skill in the art.

Chemical methods of gene therapy involve carrier mediated gene transfer through the use of fusogenic lipid



vesicles such as liposomes or other vesicles for membrane fusion. A carrier harboring a DNA of interest can be conveniently introduced into body fluids or the bloodstream and then site-specifically directed to the target organ or  
5 tissue in the body. Liposomes, for example, can be developed which are cell specific or organ specific. The foreign DNA carried by the liposome thus will be taken up by those specific cells. Injection of immunoliposomes that are targeted to a specific receptor on certain cells can be used  
10 as a convenient method of inserting the DNA into the cells bearing the receptor. Another carrier system that has been used is the asialoglycoprotein/polylysine conjugate system for carrying DNA to hepatocytes for in vivo gene transfer.

Transfected DNA may also be complexed with other kinds  
15 of carriers so that the DNA is carried to the recipient cell and then resides in the cytoplasm or in the nucleoplasm of the recipient cell. DNA can be coupled to carrier nuclear proteins in specifically engineered vesicle complexes and carried directly into the nucleus.

Carrier mediated gene transfer may also involve the use  
20 of lipid-based proteins which are not liposomes. For example, lipofectins and cytofectins are lipid-based positive ions that bind to negatively charged DNA, forming a complex that can ferry the DNA across a cell membrane.  
25 Another method of carrier mediated gene transfer involves receptor-based endocytosis. In this method, a ligand (specific to a cell surface receptor) is made to form a complex with a gene of interest and then injected into the bloodstream; target cells that have the cell surface  
30 receptor will specifically bind the ligand and transport the ligand-DNA complex into the cell.

Biological gene therapy methodologies usually employ viral vectors to insert genes into cells. The term "vector"  
as used herein in the context of biological gene therapy  
35 means a carrier that can contain or associate with specific polynucleotide sequences and which functions to transport the specific polynucleotide sequences into a cell. The

transfected cells may be cells derived from the patient's normal tissue, the patient's diseased tissue, or may be non-patient cells. Examples of vectors include plasmids and infective microorganisms such as viruses, or non-viral  
5 vectors such as the ligand-DNA conjugates, liposomes, and lipid-DNA complexes discussed above.

It may be desirable that a recombinant DNA molecule comprising a TREPA DNA sequence is operatively linked to an expression control sequence to form an expression vector  
10 capable of expressing TREPA. Alternatively, gene regulation of TREPA may be accomplished by administering proteins that bind to the kringle gene, or control regions associated with the TREPA gene, or its corresponding RNA transcript to modify the rate of transcription or translation.

15 Viral vectors that have been used for gene therapy protocols include but are not limited to, retroviruses, other RNA viruses such as poliovirus or Sindbis virus, adenovirus, adeno-associated virus, herpes viruses, SV 40, vaccinia and other DNA viruses. Replication-defective  
20 murine retroviral vectors are the most widely utilized gene transfer vectors. Murine leukemia retroviruses are composed of a single strand RNA complexed with a nuclear core protein and polymerase (pol) enzymes, encased by a protein core (gag) and surrounded by a glycoprotein envelope (env) that  
25 determines host range. The genomic structure of retroviruses include the gag, pol, and env genes enclosed by 5' and 3' long terminal repeats (LTR). Retroviral vector systems exploit the fact that a minimal vector containing the 5' and 3' LTRs and the packaging signal are sufficient  
30 to allow vector packaging, infection and integration into target cells providing that the viral structural proteins are supplied in trans in the packaging cell line.

Fundamental advantages of retroviral vectors for gene transfer include efficient infection and gene expression in  
35 most cell types, precise single copy vector integration into target cell chromosomal DNA, and ease of manipulation of the retroviral genome. For example, altered retrovirus vectors

have been used in ex vivo methods to introduce genes into peripheral and tumor-infiltrating lymphocytes, hepatocytes, epidermal cells, myocytes, or other somatic cells (which may then introduced into the patient to provide the gene product  
5 from the inserted DNA).

The adenovirus is composed of linear, double stranded DNA complexed with core proteins and surrounded with capsid proteins. Advances in molecular virology have led to the ability to exploit the biology of these organisms to create  
10 vectors capable of transducing novel genetic sequences into target cells in vivo. Adenoviral-based vectors will express gene product peptides at high levels. Adenoviral vectors have high efficiencies of infectivity, even with low titers of virus. Additionally, the virus is fully infective  
15 as a cell free virion so injection of producer cell lines is not necessary. Another potential advantage to adenoviral vectors is the ability to achieve long term expression of heterologous genes  
in vivo .

20 Viral vectors have also been used to insert genes into cells using in vivo protocols. To direct tissue-specific expression of foreign genes, cis-acting regulatory elements or promoters that are known to be tissue specific can be used. Alternatively, this can be achieved using in situ  
25 delivery of DNA or viral vectors to specific anatomical sites in vivo. A viral vector can be delivered directly to the in vivo site, by a catheter, for example, thus allowing only certain areas to be infected by the virus, and providing long-term, site-specific gene expression. In vivo  
30 gene transfer using retrovirus vectors has also been demonstrated in mammary tissue and hepatic tissue by injection of the altered virus into blood vessels leading to the organs.

When used in the above or other treatments, a  
35 therapeutically effective amount of one of the proteins of the present invention may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt form.

By a "therapeutically effective amount" of the protein of the invention is meant a sufficient amount of the protein to treat an angiogenic disease, (for example, to limit tumor growth or to slow or block tumor metastasis) at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the proteins and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific protein employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific protein employed; the duration of the treatment; drugs used in combination or coincidental with the specific protein employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the protein at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The proteins of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethansulfonate (isethionate), lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, prcrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and

undecanoate. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such  
5 inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as maleic acid, succinic acid and citric acid. Other salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic basis.  
10 Preferred salts of the proteins of the invention include phosphate, tris and acetate.

The total daily dose of the proteins of this invention administered to a human or lower animal may range from about 0.0001 to about 1 mg/kg of a patient's body mass/day. If  
15 desired, the effective daily dose may be divided into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

Alternatively, a protein of the present invention may  
20 be administered as a pharmaceutical composition containing the protein of interest in combination with one or more pharmaceutically acceptable excipients. A pharmaceutically acceptable carrier or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material  
25 or formulation auxiliary of any type. The compositions may be administered parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), rectally, or buccally. The term "parenteral" as used herein refers to  
30 modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions for parenteral injection comprise pharmaceutically-acceptable sterile aqueous or  
35 nonaqueous solutions, dispersions, suspensions or emulsions, as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use.

Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters) and poly(anhydrides). Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable mediums just prior to use.

Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical administration, including those for inhalation, may be prepared as a dry powder which may be  
5 pressurized or non-pressurized. In non-pressurized powder compositions, the active ingredient in finely divided form may be used in admixture with a larger-sized pharmaceutically-acceptable inert carrier comprising  
10 particles having a size, for example, of up to 100 micrometers in diameter. Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 micrometers.

Alternatively, the composition may be pressurized and  
15 contain a compressed gas, such as nitrogen or a liquified gas propellant. The liquified propellant medium and indeed the total composition is preferably such that the active ingredient does not dissolve therein to any substantial extent. The pressurized composition may also contain a  
20 surface active agent, such as a liquid or solid non-ionic surface active agent or may be a solid anionic surface active agent. It is preferred to use the solid anionic surface active agent in the form of a sodium salt.

A further form of topical administration is to the eye.  
25 A protein of the invention is delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the protein is maintained in contact with the ocular surface for a sufficient time period to allow the protein to penetrate the corneal and internal regions of the eye, as,  
30 for example, the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically-acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an encapsulating  
35 material. Alternatively, the protein of the invention may be injected directly into the vitreous and aqueous humour.

Compositions for rectal or vaginal administration are preferably suppositories which may be prepared by mixing the proteins of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active protein.

Proteins of the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a protein of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidylcholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

While the proteins of the invention can be administered as the sole active pharmaceutical agent, they may also be used in combination with one or more agents which are conventionally administered to patients for treating angiogenic diseases. For example, when used in the treatment of solid tumors, proteins of the invention may be administered with anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovin rescue), doxorubicin, cyclophosphamide, taxol, etoposide/mechlorethamine, vincristine, prednisone and



procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and the like.

5       Total daily dose of TREPA (administered in combination with a protein of this invention) to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily and more usually from 1 to 300 mg/kg body  
10   weight.

      It will be understood that agents which can be combined with the protein of the present invention for the inhibition, treatment or prophylaxis of angiogenic diseases are not limited to those listed above, but include in  
15   principle any agents useful for the treatment or prophylaxis of angiogenic diseases.

      Synthetic peptide fragments of TREPA may also be produced and used in a variety of applications. As examples, different peptide fragments of TREPA can be used  
20   (1) as agonists and antagonists active at TREPA binding sites, (2) as a means to isolate a TREPA receptor, (3) as antigens for the development of specific antisera, (4) as peptides for use in diagnostic kits and (5) as peptides linked to or used in combination with cytotoxic agents (for  
25   targeted killing of cells that bind TREPA). The amino acid sequences that comprise these peptides may be selected on the basis of their position on the exterior regions of the molecule which are accessible for binding to antisera. Furthermore, these peptide sequences may be compared to  
30   known sequences using protein sequence databases such as GenBank, Brookhaven Protein, SWISS-PROT, and PIR to determine potential sequence homologies. This information facilitates elimination of sequences that exhibit a high degree of sequence homology to other molecules, thereby  
35   enhancing the potential for high specificity in the development of antisera, agonists and antagonists to TREPA.

Systematic substitution of amino acids within these synthesized peptides may yield high affinity peptide agonists and antagonists to the TREPA receptor that enhance or diminish TREPA binding to its receptor. Such agonists  
5 may be used to suppress the growth of micrometastases, thereby limiting the spread of cancer. In cases of inadequate vascularization, antagonists to TREPA may be applied to block the inhibitory effects of TREPA and promote angiogenesis. For example, this type of treatment may have  
10 therapeutic effects in promoting wound healing in diabetics.

TREPA peptides may also be employed to develop affinity columns for isolation of a TREPA receptor in, for example, cultured endothelial cells. As is known in the art, isolation and purification of a TREPA receptor may be  
15 followed by amino acid sequencing to identify and isolate polynucleotides which encode the TREPA receptor. Such polynucleotides may then be cloned into a suitable expression vector and transfected into tumor cells. Expression of the receptor by the transfected tumor cells  
20 would enhance the responsiveness of these cells to endogenous or exogenous TREPA, thereby decreasing the rate of metastatic growth. Furthermore, recombinant expression of this receptor would allow greater amounts of receptor to be produced, e.g., to produce a sufficient quantity for use  
25 in high throughput screening assays to identify smaller antagonists which mimic the action of TREPA.

TREPA peptides of the present invention can also be used as antigens to generate polyclonal or monoclonal antibodies that are specific for the TREPA inhibitor. One  
30 way in which such antibodies could be used is in diagnostic methods and kits to detect or quantify TREPA in a body fluid or tissue. Results from these tests could be used to diagnose or determine the prognostic relevance of TREPA.

TREPA peptides may be chemically coupled to isotopes, 35 enzymes, carrier proteins, cytotoxic agents, fluorescent molecules, chemiluminescent, bioluminescent and other proteins for a variety of applications. For example, a

TREPA polypeptide may be labeled to facilitate testing of its ability to bind TREPA antisera or to detect cell types which possess a TREPA receptor. The coupling technique is generally chosen on the basis of the functional groups available on the amino acids of the TREPA sequence including, but not limited to amino, sulfhydryl, carboxyl, amide, phenol, and imidazole. Various reagents used to effect such couplings include among others, glutaraldehyde, diazodized benzidine, carbodiimide, and p-benzoquinone.

5  
10 The efficiency of the coupling reaction is determined using different techniques appropriate for the specific reaction. For example, radiolabeling of a TREPA peptide with  $I^{125}$  may be accomplished using chloramine T and  $NaI^{125}$  of high specific activity. The reaction is terminated with sodium metabisulfite and the mixture is desalted on  
15 disposable columns. The labeled peptide is eluted from the column, and fractions are collected. Aliquots are removed from each fraction and radioactivity measured in a gamma counter. In this manner, a labeled TREPA peptide may be  
20 obtained which is free from unreacted  $NaI^{125}$ .

Another application of peptide conjugation is for production of polyclonal antisera. For example, TREPA peptides containing lysine residues may be linked to purified bovine serum albumin using glutaraldehyde. The  
25 efficiency of this reaction may be determined by measuring the incorporation of radiolabeled peptide. Unreacted glutaraldehyde and peptide may be separated by dialysis and the conjugate stored for subsequent use.

The production of antiserum against TREPA, TREPA  
30 analogs, peptide fragments of TREPA and the TREPA receptor can be performed using established techniques known to those skilled in the art. For example, polyclonal antisera may be raised in rabbits, sheep, goats or other animals. TREPA peptides conjugated to a carrier molecule such as bovine  
35 serum albumin, or TREPA itself, may be combined with an adjuvant mixture, emulsified and injected subcutaneously at multiple sites on the back, neck, flanks, and sometimes in

the footpads of a suitable host. Generally, booster injections are then given at regular intervals, such as every 2 to 4 weeks. Approximately 7 to 10 days after each injection, blood samples are obtained by venipuncture, using, for example, the marginal ear veins after dilation. The blood samples are allowed to clot overnight at 4°C and are centrifuged at approximately 2400 X g at 4°C for about 30 minutes. The serum is removed, aliquoted, and stored at 4°C for immediate use or at -20 to -90°C for subsequent analysis.

Serum samples from generation of polyclonal antisera or media samples from production of monoclonal antisera may be analyzed for determination of antibody titer and, in particular, for the determination of high titer antisera. Subsequently, the highest titer TREPA antisera may be tested to establish the following: a) optimal antiserum dilution for highest specific binding of the antigen and lowest non-specific binding, b) ability to bind increasing amounts of TREPA peptide in a standard displacement curve, c) potential cross-reactivity with related peptides and proteins, including plasminogen and also TREPA of related species, and d) ability to detect TREPA peptides in extracts of plasma, urine, tissues, and in cell culture media.

Titer may be established through several means known in the art, such as by dot blot and density analysis, and also by precipitation of radiolabeled peptide-antibody complexes using protein A, secondary antisera, cold ethanol or charcoal-dextran followed by activity measurement with a gamma counter. If desired, the highest titer antisera may be purified on affinity columns. For example, TREPA peptides may be coupled to a commercially available resin and used to form an affinity column. Antiserum samples may then be passed through the column so that TREPA antibodies bind (via TREPA) to the column. These bound antibodies are subsequently eluted, collected and evaluated for determination of titer and specificity.

The present invention also provides a method of screening a plurality of compounds for specific binding to a TREPA derived polypeptide, or any fragment thereof, to identify at least one compound which specifically binds the TREPA derived polypeptide. Such a method comprises the steps of providing at least one compound; combining the TREPA derived polypeptide with each compound under suitable conditions for a time sufficient to allow binding; and detecting TREPA polypeptide binding to each compound.

Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the polypeptide of the present invention, is used to design an antisense RNA oligonucleotide of from 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription, thereby preventing transcription and the production of the TREPA derived polypeptide. For triple helix, see, for example, Lee et al, Nucl. Acids Res. 6:3073 (1979); Cooney et al, Science 241:456 (1988); and Dervan et al, Science 251:1360 (1991). The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of a mRNA molecule into the TREPA derived polypeptide. For antisense, see, for example, Okano, J. Neurochem. 56:560 (1991); and "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression", CRC Press, Boca Raton, Fla. (1988). Antisense oligonucleotides act with greater efficacy when modified to contain artificial internucleotide linkages which render the molecule resistant to nucleolytic cleavage. Such artificial internucleotide linkages include but are not limited to methylphosphonate, phosphorothiolate and phosphoroamidate internucleotide linkages.

The polypeptide or peptide fragment employed in such a test may either be free in solution, affixed to a solid

support, borne on a cell surface or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids which can express the polypeptide or peptide fragment. Drugs may be screened against such transformed cells in competitive binding assays. For example, the formation of complexes between a polypeptide and the agent being tested can be measured in either viable or fixed cells.

10       The present invention thus provides methods of screening for drugs or any other agent which can be used to treat diseases associated with the TREPA gene by measuring the effect of the drug on the amount of TREPA nucleotide or protein produced or biological effects of TREPA. Examples of these types of measurements include but are not limited to measuring  $\text{Ca}^{++}$  efflux, cAMP production, aptosis, etc. These measurements are known to those of ordinary skill in the art. The present invention also provides for measuring the effect of the drug on a recombinant reporter gene designed to respond to TREPA. These methods comprise measuring the effect of applying the drug to a natural or genetically engineered experimental organism, such as cultured cells, bacteria, or laboratory animals and measuring the amount of TREPA nucleotide or protein produced, or biological effects of TNF gamma, or quantity of a recombinant reporter gene such as luciferase.

      The present invention thus provides methods of screening for drugs or any other agent which can be used to treat diseases associated with the TREPA gene. These methods comprise contacting the drug with a polypeptide or fragment thereof and assaying for either the presence of a complex between the agent and the polypeptide, or for the presence of a complex between the polypeptide and the cell. In competitive binding assays, the polypeptide typically is labeled. After suitable incubation, free (or uncomplexed) polypeptide or fragment thereof is separated from that present in bound form, and the amount of free or uncomplexed

label is a measure of the ability of the particular drug to bind to polypeptide or to interfere with the polypeptide/cell complex.

The present invention also encompasses the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptide specifically compete with a test drug for binding to the polypeptide or fragment thereof. In this manner, the antibodies can be used to detect the presence of any polypeptide in the test sample which shares one or more antigenic determinants with a polypeptide provided herein.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to at least one polypeptide disclosed herein. Briefly, large numbers of different small peptide test compounds are synthesized on a solid phase, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptide and washed. Polypeptide thus bound to the solid phase is detected by methods well-known in the art. Purified polypeptide can also be coated directly onto plates for use in the drug screening techniques described herein. In addition, non-neutralizing antibodies can be used to capture the polypeptide and immobilize it on the solid support. See, for example, EP 84/03564, published on September 13, 1984, which is incorporated herein by reference

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of the small molecules including agonists, antagonists, or inhibitors with which they interact. Such structural analogs can be used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide in vivo. J. Hodgson, Bio/Technology 9:19-21 (1991), incorporated herein by reference.

For example, in one approach, the three-dimensional structure of a polypeptide, or of a polypeptide-inhibitor

complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to  
5 determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous polypeptide-like  
10 molecules or to identify efficient inhibitors

Useful examples of rational drug design may include molecules which have improved activity or stability as shown by S. Braxton et al., Biochemistry 31:7796-7801 (1992), or which act as inhibitors, agonists, or antagonists of native  
15 peptides as shown by S. B. P. Athauda et al., J. Biochem. (Tokyo) 113 (6):742-746 (1993), incorporated herein by reference.

It also is possible to isolate a target-specific antibody, selected by an assay as described hereinabove, and  
20 then to determine its crystal structure. In principle, this approach yields a pharmacophore upon which subsequent drug design can be based. It further is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies ("anti-ids") to a functional,  
25 pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-id is an analog of the original receptor. The anti-id then could be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then  
30 can act as the pharmacophore (that is, a prototype pharmaceutical drug).

A sufficient amount of a recombinant polypeptide of the present invention may be made available to perform analytical studies such as X-ray crystallography. In  
35 addition, knowledge of the polypeptide amino acid sequences which are derivable from the nucleic acid sequence provided herein will provide guidance to those employing computer



modeling techniques in place of or in addition to x-ray crystallography.

Antibodies specific to the TREPA derived polypeptide may further be used to inhibit the biological action of the polypeptide by binding to the polypeptide. In this manner, the antibodies may be used in therapy, for example, to treat TREPA diseases including inflammation.

Further, such antibodies can detect the presence or absence of TREPA derived polypeptide and, therefore, are useful as diagnostic markers for the diagnosis of TREPA disease, especially inflammation. Such antibodies may also function as diagnostic markers for TREPA disease conditions such as inflammation. The present invention also is directed to antagonists and inhibitors of the polypeptides of the present invention. The antagonists and inhibitors are those which inhibit or eliminate the function of the polypeptide. Thus, for example, an antagonist may bind to a polypeptide of the present invention and inhibit or eliminate its function. The antagonist, for example, could be an antibody against the polypeptide which eliminates the activity of TREPA-derived polypeptide by binding to TREPA-derived polypeptide, or in some cases, the antagonist may be an oligonucleotide. Examples of small molecule inhibitors include but are not limited to small peptides or peptide-like molecules.

The antagonists and inhibitors may be employed as a composition with a pharmaceutically acceptable carrier, including but not limited to saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. Administration of TREPA-derived polypeptide inhibitors are preferably systemic. The present invention also provides an antibody which inhibits the action of such polypeptides.

#### Recombinant Technology

The present invention provides host cells and expression vectors comprising polynucleotides of the present invention and methods for the production of polypeptides they encode. Such methods comprise culturing the host cells

under conditions suitable for the expression of the TREPA-derived polynucleotide and recovering the TREPA-derived polypeptide from the cell culture.

The present invention also provides vectors which  
5 include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the present invention and the production of polypeptides of the present invention by recombinant techniques.

Host cells are genetically engineered (transduced or  
10 transformed or transfected) with the vectors of this invention which may be cloning vectors or expression vectors. The vector may be in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as  
15 appropriate for activating promoters, selecting transformants or amplifying TREPA-derived genes. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled  
20 artisan.

The polynucleotide of the present invention may be employed for producing a polypeptide by recombinant techniques. Thus, the polynucleotide sequence may be included in any one of a variety of expression vehicles, in  
25 particular vectors or plasmids, for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; yeast plasmids; vectors derived from combinations of plasmids and  
30 phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudo-rabies. However, any other plasmid or vector may be used so long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the  
35 vector by a variety of procedures. In general, the DNA sequence is inserted into appropriate restriction endonuclease sites by procedures known in the art. Such

2025 RELEASE UNDER E.O. 14176

procedures and others are deemed to be within the scope of those skilled in the art. The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA  
5 synthesis. Representative examples of such promoters include but are not limited to the LTR or SV40 promoter, the E. coli lac or trp promoters, the phage lambda P sub L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.  
10 The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. In addition, the expression vectors preferably contain a gene to provide a phenotypic trait for  
15 selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

The vector containing the appropriate DNA sequence as  
20 hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein. As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Salmonella  
25 typhimurium and Streptomyces sp.; fungal cells, such as yeast; insect cells such as Drosophila and Sf9; animal cells such as CHO, COS or Bowes melanoma; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings  
30 provided herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs  
comprise a vector, such as a plasmid or viral vector, into  
35 which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory

sequences including, for example, a promoter operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available. The following vectors are provided by way of example. Bacterial: pSPORT1 (Life Technologies, Gaithersburg, MD), pQE70, pQE60, pQE-9 (Qiagen) pBs, phagescript, psiX174, pBluescript SK, pBsKS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLneo, pSV2cat, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as it is replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, SP6, T7, gpt, lambda P sub R, P sub L and trp. Eukaryotic promoters include cytomegalovirus (CMV), immediate early, herpes simplex virus (HSV) thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention provides host cells containing the above-described construct. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (L. Davis et al., "Basic Methods in Molecular Biology", 2nd edition, Appleton and Lang, Paramount Publishing, East Norwalk, CT (1994)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides

of the invention can be synthetically produced by conventional peptide synthesizers.

Proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, (Cold Spring Harbor, N.Y., 1989), which is hereby incorporated by reference.

Transcription of a DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin (bp 100 to 270), a cytomegalovirus early promoter enhancer, a polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of E. coli and S. cerevisiae TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), alpha factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting

desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and, if desirable, to provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, although others may also be employed as a routine matter of choice.

Useful expression vectors for bacterial use comprise a selectable marker and bacterial origin of replication derived from plasmids comprising genetic elements of the well-known cloning vector pBR322 (ATCC 37017). Other vectors include but are not limited to PKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is derepressed by appropriate means (e.g., temperature shift or chemical induction), and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents; such methods are well-known to the ordinary artisan.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts described by Gluzman, Cell 23:175  
5 (1981), and other cell lines capable of expressing a compatible vector, such as the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation  
10 site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the  
15 required nontranscribed genetic elements. Representative, useful vectors include pRc/CMV and pCDNA3 (available from Invitrogen, San Diego, CA).

The TREPA polypeptide is recovered and purified from recombinant cell cultures by known methods including  
20 ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, hydroxyapatite chromatography or lectin chromatography. It is preferred to have low concentrations (approximately 0.1-5  
25 mM) of calcium ion present during purification (Price, et al., J. Biol. Chem. 244:917 (1969)). Protein refolding steps can be used, as necessary, in completing configuration of the protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification  
30 steps.

The polypeptides of the present invention may be naturally purified products expressed from a high expressing cell line, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or  
35 eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production

procedure, the polypeptides of the present invention may be glycosylated with mammalian or other eukaryotic carbohydrates or may be non-glycosylated. The polypeptides of the invention may also include an initial methionine amino acid residue.

The present invention further includes modified versions of the TREPA polypeptide to preclude glycosylation while allowing expression of a reduced carbohydrate form of the protein in yeast, insect or mammalian expression systems. Known methods for inactivating glycosylation sites include, but are not limited to, those presented in U.S. patent 5,071,972 and EP 276,846, which are incorporated herein by reference.

Other variants included in the present invention include removal of sequences encoding cysteine residues, thereby preventing formation of incorrect intramolecular disulfide bridges which decrease biological activity of the protein product. The present invention also includes removal of sites of proteolytic processing, allowing expression in system which contain the problematic protease, for example, the KEX2 protease in yeast. Known methods for removing such protease sites include but are not limited to one method for removing KEX2 sites presented in EP212,914.

The present invention includes TREPA peptides in the form of oligomers, dimers, trimers and higher order oligomers. Oligomers may be formed by several means including but not limited to disulfide bonds between peptides, non-covalent interactions between peptides, and poly-ethylene-glycol linkages between peptides.

The fusion of TREPA peptides to peptide linkers or peptides that are capable of promoting oligomers is also encompassed in this invention. Such peptides include but are not limited to leucine zippers and antibody derived peptides, such as is described in Landschulz et al., Science 240:1759 (1988); Hollenbaugh and Aruffo, "Construction of Immunoglobulin Fusion Proteins", in Current Protocols in



Immunology, Supplement 4, pgs 10.19.1-10.19.11 (1992) John Wiley and sons, New York, NY.

The starting plasmids can be constructed from available plasmids in accordance with published, known procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the skilled artisan.

The following is the general procedure for the isolation and analysis of cDNA clones. In a particular embodiment disclosed herein, mRNA was isolated from TREPA and used to generate the cDNA library. TREPA was obtained from synovium of patients with rheumatoid arthritis by surgical extraction.

A cDNA insert from an isolate of TREPA was sequenced in its entirety, analyzed in detail as set forth in the Examples and is disclosed in the Sequence Listing as SEQUENCE ID NO 1. These polynucleotides encode a sufficient portion of the gene of interest to encode a biologically active molecule. The lack of full length clones is attributed to the fact that many genes are several hundred, and sometimes several thousand, bases in length and, with current technology, cannot be cloned in their entirety because of vector limitations, incomplete reverse transcription of the first strand, or incomplete replication of the second strand. Contiguous, secondary clones containing additional nucleotide sequence may be obtained using a variety of methods known to those of skill in the art.

Methods for DNA sequencing are well known in the art. Conventional enzymatic methods employ DNA polymerase, Klenow fragment, Sequenase (US Biochemical Corp, Cleveland, OH) or Taq polymerase to extend DNA chains from an oligonucleotide primer annealed to the DNA template of interest. Methods have been developed for the use of both single-stranded and double-stranded templates. The chain termination reaction products may be electrophoresed on urea/polyacrylamide gels and detected either by autoradiography (for radionucleotide labeled precursors) or by fluorescence (for fluorescent-

labeled precursors). Recent improvements in mechanized reaction preparation, sequencing and analysis using the fluorescent detection method have permitted expansion in the number of sequences that can be determined per day using machines such as the Applied Biosystems 377 DNA Sequencers (Applied Biosystems, Foster City, CA).

The reading frame of the nucleotide sequence can be ascertained by several types of analyses. First, reading frames contained within the coding sequence can be analyzed for the presence of start codon ATG and stop codons TGA, TAA or TAG. Typically, one reading frame will continue throughout the major portion of a cDNA sequence while the other two reading frames tend to contain numerous stop codons. In such cases reading frame determination is straightforward. In other more difficult cases, further analysis is required. Ultimate confirmation of a correct open reading frame is achieved by using the nucleotide sequence to produce a biologically active molecule, by such methods that are familiar to those of skill in the art.

Algorithms have been created to analyze the occurrence of individual nucleotide bases at each putative codon triplet. See, for example J. W. Fickett, Nuc Acids Res 10:5303 (1982). Coding DNA for particular organisms (bacteria, plants, and animals) tends to contain certain nucleotides within certain triplet periodicities, such as a significant preference for pyrimidines in the third codon position. These preferences have been incorporated into widely available software which can be used to determine coding potential (and frame) of a given stretch of DNA. The algorithm-derived information combined with start/stop codon information can be used to determine proper frame with a high degree of certainty. This, in turn, readily permits cloning of the sequence in the correct reading frame into appropriate expression vectors.

The nucleic acid sequences disclosed herein may be joined to a variety of other polynucleotide sequences and vectors of interest by means of well established recombinant

DNA techniques. See J. Sambrook et al., supra. Vectors of interest include cloning vectors such as plasmids, cosmids, phage derivatives, phagemids, as well as sequencing, replication, and expression vectors, and the like. In  
5 general, such vectors contain an origin of replication functional in at least one organism, convenient restriction endonuclease digestion sites, and selectable markers appropriate for particular host cells. The vectors can be transferred by a variety of means known to those of skill in  
10 the art into suitable host cells which then produce the desired DNA, RNA or polypeptides.

Occasionally, sequencing or random reverse transcription errors will mask the presence of the appropriate open reading frame or regulatory element. In  
15 such cases, it is possible to determine the correct reading frame by attempting to express the polypeptide and determining the amino acid sequence by standard peptide mapping and sequencing techniques. See, F. M. Ausubel, et al., Current Protocols in Molecular Biology, John Wiley &  
20 Sons, New York, NY (1989). Additionally, the actual reading frame of a given nucleotide sequence may be determined by transfection of host cells with vectors containing all three potential reading frames. Only those cells with the nucleotide sequence in the correct reading frame will  
25 produce a peptide of the predicted length.

The nucleotide sequences provided herein have been prepared by current, state-of-the-art, automated methods and as such may contain unidentified nucleotides. These will not present a problem to those skilled in the art who wish  
30 to practice the invention. Several methods employing standard recombinant techniques, described in J. Sambrook (supra) or periodic updates thereof, may be used to complete the missing sequence information. The same techniques used for obtaining a full length sequence, as described herein,  
35 may be used to obtain nucleotide sequence.

Expression of a particular cDNA may be accomplished by subcloning the cDNA into an appropriate expression vector

and transfecting this vector into an appropriate expression host. The cloning vector used for the generation of the TREPA cDNA library can be used for transcribing mRNA of a particular cDNA. Immediately following these eight residues  
 5 is an engineered bacteriophage promoter useful for artificial priming and transcription and a number of unique restriction sites, including EcoR I, for cloning. The vector can be transfected into an appropriate host strain of E. coli.

10 Induction of the isolated bacterial strain with isopropylthiogalactoside (IPTG) using standard methods will produce a fusion protein which contains the first seven residues of beta-galactosidase, about 15 residues of linker, and the peptide encoded within the cDNA. Since cDNA clone  
 15 inserts are generated by an essentially random process, there is one chance in three that the included cDNA will lie in the correct frame for proper translation. If the cDNA is not in the proper reading frame, the correct frame can be obtained by deletion or insertion of an appropriate number  
 20 of bases by well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or oligonucleotide linker inclusion.

The cDNA can be shuttled into other vectors known to be useful for expression of protein in specific hosts.  
 25 Oligonucleotide primers containing cloning sites and segments of DNA sufficient to hybridize to stretches at both ends of the target cDNA can be synthesized chemically by standard methods. These primers can then be used to amplify the desired gene segments by PCR. The resulting new gene  
 30 segments can be digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments can be produced by digestion of the cDNA with appropriate restriction enzymes and filling in the missing gene segments  
 35 with chemically synthesized oligonucleotides. Segments of the coding sequence from more than one gene can be ligated

together and cloned in appropriate vectors to optimize expression of recombinant sequence.

Suitable expression hosts for such chimeric molecules include but are not limited to mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells, insect cells such as Sf9 cells, yeast cells such as Saccharomyces cerevisiae, and bacteria such as E. coli. For each of these cell systems, a useful expression vector may also include an origin of replication to allow propagation in bacteria and a selectable marker such as the beta-lactamase antibiotic resistance gene to allow selection in bacteria. In addition, the vectors may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts may require the addition of 3' poly A tail if the sequence of interest lacks poly A, and/or the addition of an intron sequence which promotes proper splicing and processing of the mRNA, but does not alter the amino acid sequence of the gene product.

Further, this invention encompasses expression vectors whereby secretion of the protein outside of the host cell is achieved by fusing in frame DNA encoding signal peptide sequences to the open reading frame encoding TREPA. These sequences may be prokaryotic, eukaryotic, or viral in origin.

Additionally, the vector may contain promoters or enhancers which increase gene expression. Such promoters are host specific and include but are not limited to MMTV, SV40, or metallothionine promoters for CHO cells; trp, lac, tac or T7 promoters for bacterial hosts; or alpha factor, alcohol oxidase or PGH promoters for yeast. Adenoviral vectors with or without transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to drive protein expression in mammalian cell lines. Once homogeneous cultures of recombinant cells are obtained, large quantities of recombinantly produced protein can be recovered from the conditioned medium and analyzed using

chromatographic methods well known in the art. An alternative method for the production of large amounts of secreted protein involves the transformation of mammalian embryos and the recovery of the recombinant protein from milk produced by transgenic cows, goats, sheep, etc. Polypeptides and closely related molecules may be expressed recombinantly in such a way as to facilitate protein purification. One approach involves expression of a chimeric protein which includes one or more additional polypeptide domains not naturally present on human polypeptides. Such purification-facilitating domains include, but are not limited to, metal-chelating peptides such as histidine-tryptophan domains that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle, WA). The inclusion of a cleavable linker sequence such as Factor XA or enterokinase from Invitrogen (San Diego, CA) between the polypeptide sequence and the purification domain may be useful for recovering the polypeptide.

#### Immunoassays

The polypeptides including their fragments or derivatives or analogs thereof of the present invention, or cells expressing them, can be used in a variety of assays, many of which are described herein, for the detection of antibodies to TREPA. They also can be used as an immunogen to produce antibodies. These antibodies can be, for example, polyclonal or monoclonal antibodies, chimeric, single chain and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

For example, antibodies generated against a polypeptide corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptide into an animal or by administering the polypeptide to an animal such

as a mouse, rabbit, goat or human. A mouse, rabbit or goat is preferred. The antibody so obtained then will bind the polypeptide itself. In this manner, even a sequence encoding only a fragment of the polypeptide can be used to generate antibodies that bind the native polypeptide. Such antibodies can then be used to isolate the polypeptide from test samples such as tissue suspected of containing that polypeptide. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique as described by Kohler and Milstein, Nature 256:495-497 (1975), the trioma technique, the human B-cell hybridoma technique as described by Kozbor et al, Immun. Today 4:72 (1983), and the EBV-hybridoma technique to produce human monoclonal antibodies as described by Cole, et al., in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc, New York, NY, pp. 77-96 (1985). Techniques described for the production of single chain antibodies can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. See, for example, U.S. Pat. No. 4,946,778, which is incorporated herein by reference.

Various assay formats may utilize the antibodies of the present invention, including "sandwich" immunoassays and probe assays. For example, the monoclonal antibodies or fragments thereof of the present invention can be employed in various assay systems to determine the presence, if any, of TREPA derived polypeptide in a test sample. For example, in a first assay format, a polyclonal or monoclonal antibody or fragment thereof, or a combination of these antibodies, which has been coated on a solid phase, is contacted with a test sample, to form a first mixture. This first mixture is incubated for a time and under conditions sufficient to form antigen/antibody complexes. Then, an indicator reagent comprising a monoclonal or a polyclonal antibody or a fragment thereof, or a combination of these antibodies, to which a signal generating compound has been attached, is

contacted with the antigen/antibody complexes to form a second mixture. This second mixture then is incubated for a time and under conditions sufficient to form antibody/antigen/antibody complexes. The presence of a  
5 TREPA-derived polypeptide antigen present in the test sample and captured on the solid phase, if any, is determined by detecting the measurable signal generated by the signal generating compound. The amount of TREPA derived polypeptide antigen present in the test sample is  
10 proportional to the signal generated.

Or, a polyclonal or monoclonal TREPA-derived polypeptide antibody or fragment thereof, or a combination of these antibodies which is bound to a solid support, the test sample and an indicator reagent comprising a monoclonal  
15 or polyclonal antibody or fragments thereof, which specifically binds to TREPA derived polypeptide antigen, or a combination of these antibodies to which a signal generating compound is attached, are contacted to form a mixture. This mixture is incubated for a time and under  
20 conditions sufficient to form antibody/antigen/antibody complexes. The presence, if any, of TREPA-derived polypeptide present in the test sample and captured on the solid phase is determined by detecting the measurable signal generated by the signal generating compound. The amount of  
25 TREPA derived polypeptide proteins present in the test sample is proportional to the signal generated.

In another assay format, one or a combination of at least two monoclonal antibodies of the invention can be employed as a competitive probe for the detection of  
30 antibodies to TREPA-derived polypeptide protein. For example, TREPA-derived polypeptide proteins such as the recombinant antigens disclosed herein, either alone or in combination, are coated on a solid phase. A test sample suspected of containing antibody to TREPA-derived  
35 polypeptide antigen then is incubated with an indicator reagent comprising a signal generating compound and at least one monoclonal antibody of the invention for a time and



under conditions sufficient to form antigen/antibody complexes of either the test sample and indicator reagent bound to the solid phase or the indicator reagent bound to the solid phase. The reduction in binding of the monoclonal antibody to the solid phase can be quantitatively measured.

In yet another detection method, each of the monoclonal or polyclonal antibodies of the present invention can be employed in the detection of TREPA-derived polypeptide antigens in fixed tissue sections, as well as fixed cells by immunohistochemical analysis. Cytochemical analysis wherein these antibodies are labeled directly (with, for example, fluorescein, colloidal gold, horseradish peroxidase, alkaline phosphatase, etc.) or are labeled by using secondary labeled anti-species antibodies (with various labels as exemplified herein) to track the histopathology of disease also are within the scope of the present invention.

In addition, these monoclonal antibodies can be bound to matrices similar to CNBr-activated Sepharose and used for the affinity purification of specific TREPA-derived polypeptide proteins from cell cultures or biological tissues such as to purify recombinant and native TREPA-derived polypeptide antigens and proteins.

The monoclonal antibodies of the invention can also be used for the generation of chimeric antibodies for therapeutic use or other similar applications.

The monoclonal antibodies or fragments thereof can be provided individually to detect TREPA-derived polypeptide antigens. Combinations of the monoclonal antibodies (and fragments thereof) provided herein also may be used together as components in a mixture or "cocktail" of at least one TREPA-derived polypeptide antibody of the invention with antibodies to other TREPA-derived polypeptide regions, each having different binding specificities. Thus, this cocktail can include the monoclonal antibodies of the invention which are directed to TREPA-derived polypeptide proteins of TREPAS and other monoclonal antibodies to other antigenic determinants of TREPA-derived polypeptide genome.

The polyclonal antibody or fragment thereof which can be used in the assay formats should specifically bind to a TREPA-derived polypeptide region or other TREPA-derived polypeptide proteins used in the assay. The polyclonal antibody used preferably is of mammalian origin; human, goat, rabbit or sheep anti-TREPA-derived polypeptide polyclonal antibody can be used. Most preferably, the polyclonal antibody is rabbit polyclonal anti-TREPA-derived polypeptide antibody. The polyclonal antibodies used in the assays can be used either alone or as a cocktail of polyclonal antibodies. Since the cocktails used in the assay formats are comprised of either monoclonal antibodies or polyclonal antibodies having different TREPA-derived polypeptide specificity, they would be useful for diagnosis, evaluation and prognosis of TREPA-derived polypeptide conditions, as well as for studying TREPA-derived polypeptide protein differentiation and specificity.

It is contemplated and within the scope of the present invention that the TREPA-derived polypeptide may be detectable in assays by use of a recombinant antigen as well as by use of a synthetic peptide or purified peptide, which contains an amino acid sequence of TREPA-derived polypeptide. It also is within the scope of the present invention that different synthetic, recombinant or purified peptides identifying different epitopes of the TREPA-derived polypeptide can be used in combination in an assay to diagnose, evaluate, or prognosticate the TREPA disease condition. In this case, these peptides can be coated onto one solid phase, or each separate peptide may be coated on separate solid phases, such as microparticles, and then combined to form a mixture of peptides which can be later used in assays. Furthermore, it is contemplated that multiple peptides which define epitopes from different polypeptides may be used in combination to make a diagnosis, evaluation, or prognosis of TREPA disease. Peptides coated on solid phases or labelled with detectable labels are then allowed to compete with peptides from a patient sample for a

limited amount of antibody. A reduction in binding of the synthetic, recombinant, or purified peptides to the antibody (or antibodies) is an indication of the presence of TREPA-secreted polypeptides in the patient sample which in turn indicates the presence of TREPA gene in the patient. Such variations of assay formats are known to those of ordinary skill in the art and are discussed herein below.

In another assay format, the presence of antibody and/or antigen to TREPA-derived polypeptide can be detected in a simultaneous assay, as follows. A test sample is simultaneously contacted with a capture reagent of a first analyte, wherein said capture reagent comprises a first binding member specific for a first analyte attached to a solid phase and a capture reagent for a second analyte, wherein said capture reagent comprises a first binding member for a second analyte attached to a second solid phase, to thereby form a mixture. This mixture is incubated for a time and under conditions sufficient to form capture reagent/first analyte and capture reagent/second analyte complexes. These so-formed complexes then are contacted with an indicator reagent comprising a member of a binding pair specific for the first analyte labeled with a signal generating compound and an indicator reagent comprising a member of a binding pair specific for the second analyte labeled with a signal generating compound to form a second mixture. This second mixture is incubated for a time and under conditions sufficient to form capture reagent/first analyte/indicator reagent complexes and capture reagent/second analyte/indicator reagent complexes. The presence of one or more analytes is determined by detecting a signal generated in connection with the complexes formed on either or both solid phases as an indication of the presence of one or more analytes in the test sample. In this assay format, recombinant antigens derived from human expression systems may be utilized as well as monoclonal antibodies produced from the proteins derived from the mammalian expression systems as disclosed herein. Such

assay systems are described in greater detail in EP  
Publication No. 0473065.

In yet other assay formats, the polypeptides disclosed  
herein may be utilized to detect the presence of anti-TREPA  
5 derived polypeptide in test samples. For example, a test  
sample is incubated with a solid phase to which at least one  
recombinant protein has been attached. These are reacted  
for a time and under conditions sufficient to form  
antigen/antibody complexes. Following incubation, the  
10 antigen/antibody complex is detected. Indicator reagents  
may be used to facilitate detection, depending upon the  
assay system chosen. In another assay format, a test sample  
is contacted with a solid phase to which a recombinant  
protein produced as described herein is attached and also is  
15 contacted with a monoclonal or polyclonal antibody specific  
for the protein, which preferably has been labeled with an  
indicator reagent. After incubation, for a time and under  
conditions sufficient for antibody/antigen complexes to  
form, the solid phase is separated from the free phase, and  
20 the label is detected in either the solid or free phase as  
an indication of the presence of TREPA-derived polypeptide  
antibody. Other assay formats utilizing the recombinant  
antigens disclosed herein are contemplated. These include  
contacting a test sample with a solid phase to which at  
25 least one antigen from a first source has been attached,  
incubating the solid phase and test sample for a time and  
under conditions sufficient to form antigen/antibody  
complexes, and then contacting the solid phase with a  
labeled antigen, which antigen is derived from a second  
30 source different from the first source. For example, a  
recombinant protein derived from a first source such as E.  
coli is used as a capture antigen on a solid phase, a test  
sample is added to the so-prepared solid phase, and a  
recombinant protein derived from a different source (i.e.,  
35 non-E. coli) is utilized as a part of an indicator reagent.  
Likewise, combinations of a recombinant antigen on a solid  
phase and synthetic peptide in the indicator phase also are

possible. Any assay format which utilizes an antigen specific for TREPA-derived polypeptide from a first source as the capture antigen and an antigen specific for TREPA-derived polypeptide from a different second source are contemplated. Thus, various combinations of recombinant antigens, as well as the use of synthetic peptides, purified proteins, and the like, are within the scope of this invention. Assays such as this and others are described in U.S. Patent No. 5,254,458, which enjoys common ownership and is incorporated herein by reference.

Other embodiments which utilize various other solid phases also are contemplated and are within the scope of this invention. For example, ion capture procedures for immobilizing an immobilizable reaction complex with a negatively charged polymer (described in EP publication 0326100 and EP publication No. 0406473), can be employed according to the present invention to effect a fast solution-phase immunochemical reaction. An immobilizable immune complex is separated from the rest of the reaction mixture by ionic interactions between the negatively charged poly-anion/immune complex and the previously treated, positively charged porous matrix and detected by using various signal generating systems previously described, including those described in chemiluminescent signal measurements as described in EPO Publication No. 0 273,115.

Also, the methods of the present invention can be adapted for use in systems which utilize microparticle technology including in automated and semi-automated systems wherein the solid phase comprises a microparticle (magnetic or non-magnetic). Such systems include those described in published EPO applications Nos. EP 0 425 633 and EP 0 424 634, respectively.

The use of scanning probe microscopy (SPM) for immunoassays also is a technology to which the monoclonal antibodies of the present invention are easily adaptable. In scanning probe microscopy, in particular, in atomic force microscopy, the capture phase, for example, at least one of

the monoclonal antibodies of the invention, is adhered to a solid phase and a scanning probe microscope is utilized to detect antigen/antibody complexes which may be present on the surface of the solid phase. The use of scanning

5 tunneling microscopy eliminates the need for labels which normally must be utilized in many immunoassay systems to detect antigen/antibody complexes. The use of SPM to monitor specific binding reactions can occur in many ways. In one embodiment, one member of a specific binding partner

10 (analyte specific substance which is the monoclonal antibody of the invention) is attached to a surface suitable for scanning. The attachment of the analyte specific substance may be by adsorption to a test piece which comprises a solid phase of a plastic or metal surface, following methods known

15 to those of ordinary skill in the art. Or, covalent attachment of a specific binding partner (analyte specific substance) to a test piece which test piece comprises a solid phase of derivatized plastic, metal, silicon, or glass may be utilized. Covalent attachment methods are known to

20 those skilled in the art and include a variety of means to irreversibly link specific binding partners to the test piece. If the test piece is silicon or glass, the surface must be activated prior to attaching the specific binding partner. Also, polyelectrolyte interactions may be used to

25 immobilize a specific binding partner on a surface of a test piece by using techniques and chemistries. The preferred method of attachment is by covalent means. Following attachment of a specific binding member, the surface may be further treated with materials such as serum, proteins, or

30 other blocking agents to minimize non-specific binding. The surface also may be scanned either at the site of manufacture or point of use to verify its suitability for assay purposes. The scanning process is not anticipated to alter the specific binding properties of the test piece.

35 While the present invention discloses the preference for the use of solid phases, it is contemplated that the reagents such as antibodies, proteins and peptides of the

present invention can be utilized in non-solid phase assay systems. These assay systems are known to those skilled in the art, and are considered to be within the scope of the present invention.

5           It is contemplated that the reagent employed for the assay can be provided in the form of a test kit with one or more containers such as vials or bottles, with each container containing a separate reagent such as a probe, primer, monoclonal antibody or a cocktail of monoclonal  
10 antibodies, or a polypeptide (either recombinant or synthetic) employed in the assay. Other components such as buffers, controls, and the like, known to those of ordinary skill in art, may be included in such test kits. It also is contemplated to provide test kits which have means for  
15 collecting test samples comprising accessible body fluids, e.g., blood, urine, saliva, and stool. Such tools useful for collection ("collection materials") include lancets and absorbent paper or cloth for collecting and stabilizing blood; swabs for collecting and stabilizing saliva; cups for  
20 collecting and stabilizing urine or stool samples. Collection materials, papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. The collection materials also may be treated with or contain preservatives,  
25 stabilizers or antimicrobial agents to help maintain the integrity of the specimens. Test kits designed for the collection, stabilization, and preservation of test specimens obtained by surgery or needle biopsy are also useful. It is contemplated that all kits may be configured  
30 in two components which can be provided separately; one component for collection and transport of the specimen, and the other component for the analysis of the specimen. The collection component, for example, can be provided to the open market user while the components for analysis can be  
35 provided to others such as laboratory personnel for determination of the presence, absence or amount of analyte. Further, kits for the collection, stabilization, and

preservation of test specimens may be configured for use by untrained personnel and may be available in the open market for use at home with subsequent transportation to a laboratory for analysis of the test sample.

5        E. coli bacteria (clone 1235095) has been deposited at the American Type Culture Collection (A.T.C.C.), 12301 Parklawn Drive, Rockville, Maryland 20852, as of September 26, 1996, under the terms of the Budapest Treaty and will be maintained for a period of thirty (30) years from the date  
10 of deposit, or for five (5) years after the last request for the deposit, or for the enforceable period of the U.S. patent, whichever is longer. The deposit and any other deposited material described herein are provided for convenience only, and are not required to practice the  
15 present invention in view of the teachings provided herein. The cDNA sequence in all of the deposited material is incorporated herein by reference. Clone 1235095 was accorded A.T.C.C. Deposit No 98184.

20        The present invention will now be described by way of examples, which are meant to illustrate, but not to limit, the scope of the present invention.

#### EXAMPLES

##### Example 1: Identification of TREPA Library EST Clones

25        A. Library Comparison of Expressed Sequence Tags (ESTs) or Transcript Images. Partial sequences of cDNA clone inserts, so-called expressed sequence tags (ESTs), were derived from cDNA libraries made from numerous tissues, both diseased and normal, and entered into a database  
30 (LIFESEQ™ database, available from Incyte Pharmaceuticals, Palo Alto, CA) as gene transcript images. The sequences then were evaluated to identify EST sequences whose predicted amino acid sequences were representative of the TNF family of genes. Once such an EST was identified, the  
35 known sequence of the gene represented by the EST was expanded by searching all available human DNA sequence databases, both publicly and privately, available for



overlapping sequences with significant homology. This expanded sequence will from here on be described as the "contig". Having done this, the 5' end furthest extending clone available at the time was obtained from the EST provider (Incyte Pharmaceuticals). None of these overlapping EST sequences in either the Incyte or publicly available databases were annotated by the producers of the sequence as being members of the TNF family of ligands. Also, no one of these sequences is sufficient to produce a soluble, active form of the protein.

#### Example 2: Sequencing of EST-Containing Clones

DNA sequences for clones which comprise the most upstream ESTs of the contig are determined using dideoxy termination sequencing with either dye-labelled primers, dye terminators, or radiolabeled nucleotides, following known methods. See, for example, F. Sanger et al., Proc. Natl. Acad. Sci. U.S.A. 74:5463 (1977).

Because the vector pSPORT1 (Life Technologies, Gaithersburg, MD) and pINCY, based on pSPORT-1, contains universal priming sites just adjacent to the 3' and 5' ligation junctions of the inserts, the inserts are sequenced in both directions using universal primers. The sequencing reactions are run on a polyacrylamide denaturing gel, and the sequences are determined by an Applied Biosystems 377 Sequencer (available from Applied Biosystems, Foster City, CA) or other sequencing apparatus.

Based upon other members of the TNF family of ligands, the sequence of clone ID #690050 (SEQUENCE ID NO1) contains the entire TNF extracellular homology domain, and is therefore sufficient to form a biologically active molecule, one example of which is SEQUENCE ID NO 3.

#### Example 3: Nucleic Acid Preparation

A. RNA Extraction from Tissue. Total RNA is isolated from solid tissues or cells from patients with prostate cancer and non-tumor tissues using a lithium chloride/urea technique known in the art and described by N. Kato et al.,

J. Virology 61:2182-2191 (1987). Non-tumor tissues are used as negative controls. The mRNA can be further purified from total RNA using commercially available kits such as oligo dT cellulose spin columns (RediCol™ from Pharmacia, Uppsala, Sweden) for the isolation of poly-adenylated RNA. Total RNA or mRNA then is dissolved in lysis buffer (5M guanidine thiocyanate, 0.1M EDTA, pH 7.0) for analysis in the ribonuclease protection assay.

B. RNA Extraction from Blood. RNA is prepared from blood samples from patients with or without diagnosed TREPA-associated disease by the standard QIAamp (Qiagen, Chatsworth, CA) RNA protocol. Briefly, 25 µl of blood are mixed with 280 µl of Qiagen AVL buffer and incubated at room temperature for 15 min. Then, 280 µl of 100% ethanol is added to the mixture, and the entire mixture is transferred to a QIAamp spin column. Next, the column is spun at 6,000 x g for 2 min, washed twice with 500 µl of Qiagen AW/ethanol buffer and spun at 6,000 x g for 2 min. The column is spun an additional 3 min at >10,000 x g. The RNA is eluted by adding 100 µl of RNase-free water preheated at 80°C to the column and spinning at 6,000 x g for 2 min.

C. RNA Extraction from polysomes. Tissue is minced in saline at 4°C and mixed with 2.5 volumes of 0.8 M sucrose in a TK150M (150 mM KCl, 5 mM MgCl<sub>2</sub>, 50 mM Tris-HCl, pH 7.4) solution containing 6 mM 2-mercaptoethanol. The tissue is homogenized in a Teflon-glass Potter homogenizer with five strokes at 100-200 rpm followed by six strokes in a Dounce homogenizer, as described by B. Mechler, Methods in Enzymology 152:241-248 (1987). The homogenate then is centrifuged at 12,000 x g for 15 min at 4°C to sediment the nuclei. The polysomes are isolated by mixing 2 ml of the supernatant with 6 ml of 2.5 M sucrose in TK150M and layering this mixture over 4 ml of 2.5 M sucrose in TK150M in a 38 ml polyallomer tube. Two additional sucrose TK150M solutions are successively layered onto the extract fraction; a first layer of 13 ml of 2.05 M sucrose followed by a second layer of 6 ml of 1.3 M sucrose. The polysomes

are isolated by centrifuging the gradient at 90,000 x g for 5 h at 4°C. The fraction then is taken from the 1.3 M sucrose/2.05 M sucrose interface with a siliconized pasteur pipette and diluted in an equal volume of TE (10 mM Tris-HCl, pH 7.4, 1 mM EDTA). An equal volume of 90°C SDS buffer (1% SDS, 200 mM NaCl, 20 mM Tris-HCl, pH 7.4) is added, and the solution is incubated in a boiling water bath for 2 min. Proteins next are digested with a proteinase-K digestion (50 mg/ml) for 15 min at 37°C. The mRNA is purified with 3 equal volumes of phenol-chloroform extractions followed by precipitation with 0.1 volume of 2 M sodium acetate (pH 5.2) and 2 volumes of 100% ethanol at -20°C overnight. The precipitated RNA is recovered by centrifugation at 12,000 x g for 10 min at 4°C. The RNA is dried and resuspended in TE, pH 7.4 or distilled water. The resuspended RNA then can be used in a slot blot or dot blot hybridization assay to check for the presence of mRNA containing EST sequences (see Example 6).

The quality of nucleic acid and proteins is dependent on the method of preparation used. Each sample may require a different preparation technique to maximize isolation efficiency of the target molecule.

#### Example 4: Ribonuclease Protection Assay

A. Labeling of Complementary RNA (cRNA) Hybridization Probes. Labeled sense and antisense riboprobes are transcribed from the EST sequence which contains a 5' RNA polymerase promoter such as SP6 or T7. The sequence may be from a vector containing the appropriate EST insert or from a PCR-generated product of the insert using PCR primers which incorporate a 5' RNA polymerase promoter sequence. The transcripts are prepared in a 20 µl reaction volume containing 1 µg of DNA template, 2 µl of 100 mM dithiothreitol, 0.8 µl of RNasin (10-40U), 500 µM each of ATP, CTP, GTP, 5 µl (alpha<sup>32</sup>P) UTP or 100-500 µM biotinylated UTP, and 1 µl of RNA polymerase in transcription buffer (40 mM Tris-HCl, pH 7.5, 6 mM MgCl<sub>2</sub>, 2 mM spermidine HCl, 5 mM NaCl). Following incubation at 37°C

for one hour, the transcripts are treated with DNase I (15 U) for an additional 30 min to digest the template. The probes then are isolated by spin columns, salt precipitation or electrophoresis techniques which are well-known in the art. Finally, the probes are dissolved in lysis buffer (5 M Guanidine Thiocyanate, 0.1 M EDTA, pH 7.0).

B. Hybridization of Labeled Probe to Target.

Approximately 20 µg of extracted total cellular RNA, as obtained in Example 3 supra, in 10 µl of lysis buffer are mixed with either (i)  $1 \times 10^5$  cpm of radioactively labeled probe or (ii) 250 pg of non-isotopically labeled probe, each in 2 µl of lysis buffer. The mixture then is incubated at 60°C for 5 min and hybridized overnight at room temperature. See, T. Kaabache et al., Anal. Biochem. 232:225-230 (1995).

C. RNase Digestion. Hybridizations are terminated by incubation with 380 µl of a solution containing 40 µg/ml RNase A and 625 units/ml RNase T1 in 1 mM EDTA, 300 mM NaCl, 30 mM Tris-HCl pH 7.4 for 45-60 min at room temperature. RNase digestion then is terminated by the addition of 60 µl of proteinase-K (1.7 mg/ml) containing 3.3% SDS, followed by incubation for 30 min at 37°C. The digested mixture then is extracted with phenol:chloroform:isoamyl alcohol to remove protein. The mRNA:cRNA hybrids are precipitated from the aqueous phase by the addition of 4 µg yeast tRNA and 800 µl of ethanol, and incubation at -80°C for 30 min. The precipitates are collected by centrifugation.

D. Fragment Analysis. The precipitates are dissolved in 5 µl of denaturing gel loading dye (80% formamide, 10 mM EDTA, pH 8.0, 1 mg/ml xylene cyanol, 1 mg/ml bromophenol blue) and electrophoresed in 6 % polyacrylamide TBE, 8 M urea denaturing gels. The gels are dried under vacuum and autoradiographed. Quantitation can be performed by comparing the counts obtained from the test samples to a calibration curve that was generating by utilizing calibrators that are the sense strand. In cases where non-isotopic labels are used, hybrids are transferred from the gels to membranes (nylon or nitrocellulose) by blotting and

then analyzed using detection systems that employ streptavidin alkaline phosphatase conjugates and chemiluminescence or chemifluorescence reagents. High level of expression of mRNA corresponding to a sequence selected from the group consisting of SEQUENCE ID NO 1, or fragments or complements thereof, then is an indication of the presence of TREPA gene.

#### Example 5: Northern Blotting

The northern blot technique is used to identify a RNA fragment of a specific size from a complex population of RNA using gel electrophoresis and nucleic acid hybridization. Northern blotting is a well-known technique in the art. Briefly, up to 20 µg of extracted RNA (see Example 3) are incubated in 20 µl of a solution containing 40 mM morpholinopropanesulfonic acid (MOPS), pH 7.0, 10 mM sodium acetate, 1 mM EDTA, 2.2 M formaldehyde, 50% v/v formamide for 15 min at 55°C. The denatured RNA is mixed with 2 µl of loading buffer (50% glycerol, 1 mM EDTA, 0.4% bromophenol blue, 0.4% xylene cyanol) and loaded into a denaturing 1.5% agarose gel containing 40 mM morpholinopropanesulfonic acid (MOPS), pH 7.0, 10 mM sodium acetate, 1 mM EDTA and 2.2 M formaldehyde. The gel is electrophoresed for an appropriate time, transferred to a wash tray and washed with five changes of RNase free water for 5 min followed by a 45 min soak at room temperature in 50 mM NaOH and 10 mM NaCl. The gel is neutralized by soaking for 45 min in 0.1 M Tris-HCl, pH 7.5. After a 1 h soak in 20X SSC buffer (3 M NaCl, 300 mM tri-sodium citrate), the gel is transferred onto a nitrocellulose or nylon based matrix. After transfer is complete, the filter is washed in 3X SSC, air dried for 2 h and baked at 80°C for 4 h under vacuum. The mRNAs are detected as in Example 4, supra. Again, high level of expression of mRNA corresponding a sequence selected from the group consisting of SEQUENCE ID NO 1 or fragments or complements thereof, is an indication of the presence of the TREPA gene.

Example 6: Dot Blot/Slot Blot

Dot and slot blot assays are quick methods to evaluate the presence of a specific nucleic acid sequence in a complex mix of nucleic acids.

- 5 To perform, up to 20  $\mu$ g of RNA is mixed in 50  $\mu$ l of 50% formamide, 7% formaldehyde, 1X SSC, incubated 15 min at 68°C and cooled on ice. Then, 100  $\mu$ l of 20X SSC is added to the RNA mixture and loaded under vacuum onto a manifold apparatus that has a prepared nitrocellulose or nylon
- 10 membrane. The membrane is soaked in water, 20X SSC for 1 hour, placed on two sheets of 20X SSC prewet Whatman #3 filter paper, and loaded into a slot blot or dot blot vacuum manifold apparatus. The slot blot is analyzed with probes prepared and labeled as in the Example 4 supra.
- 15 Detection of mRNA corresponding to a sequence selected from the group consisting of SEQUENCE ID NO 1, or fragments or complements thereof, is an indication of the presence of the TREPA gene, suggesting the diagnosis of inflammatory disease.
- 20 Other methods and buffers not specifically detailed for Examples 5 and 6 are described in J. Sambrook et al, supra.

Example 7: In Situ Hybridization

- 25 This method is useful to directly detect specific target nucleic acid sequences in cells using detectable nucleic acid hybridization probes.

- Tissues are prepared with cross-linking fixatives agents such as paraformaldehyde or glutaraldehyde for maximum cellular RNA retention. See, L. Angerer et al.,
- 30 Methods in Cell Biol. 35:37-71 (1991). Briefly, the tissue is placed in greater than 5 volumes of 1% glutaraldehyde in 50 mM sodium phosphate, pH 7.5 at 4°C for 30 min. The solution is changed with fresh solution for a further 30 min fixing. The fixing solution should have an osmolality of
- 35 approximately 0.375% NaCl. The tissue is washed once in isotonic NaCl to remove the phosphate.

The fixed tissues then are embedded in paraffin, as follows. The tissue is dehydrated through a series of ethanol

concentrations for 15 min each: 50% twice, 70% twice, 85%, 90% and 100% twice. The tissue next is soaked in two changes of xylene for 20 min each at room temperature; then it is soaked in two changes of 1 xylene:1 paraffin for 20 min each  
 5 at 60°C; and then it is soaked in three final changes in paraffin for 15 min each.

The tissue next is cut in 5  $\mu$ m sections using a standard microtome and placed on a slide previously treated with the tissue adhesive 3-aminopropyltriethoxysilane.

10 Paraffin is removed from the tissue by two 10 min xylene soaks and rehydrated in a series of ethanol concentrations; 99% twice, 95%, 85%, 70%, 50%, 30% and distilled water twice. The sections are pre-treated with 0.2 M HCl for 10 min and permeabilized with 2  $\mu$ g/ml  
 15 Proteinase-K at 37°C for 15 min.

Labeled riboprobes transcribed from the EST pSPORT1 plasmid (see Example 4) are hybridized to the prepared tissue sections and hybridized overnight at 56°C in 3X standard saline extract and 50% formamide. Excess probe is  
 20 removed by washing in 2X standard saline citrate and 50% formamide followed by digestion with 100  $\mu$ g/ml RNase A at 37°C for 30 min. Fluorescence probe is visualized by illumination with UV light under a microscope. Fluorescence in the cytoplasm is indicative of mRNA production.  
 25 Fluorescence in the nucleus detects the presence of genomic material. Alternatively, the sections can be visualized by autoradiography.

#### Example 8: Reverse Transcription PCR

30 A. One Step RT-PCR Assay. Target-specific primers are designed to detect the above target sequence by reverse transcription PCR by methods known in the art. One step RT-PCR is a sequential procedure that performs both RT and PCR in a single reaction mixture. The procedure is performed in  
 35 a 200  $\mu$ l reaction mixture containing 50 mM (N,N,-bis[2-Hydroxyethyl]glycine), pH 8.15, 81.7 mM KOAc, 33.33 mM KOH, 0.01 mg/ml bovine serum albumin, 0.1 mM ethylene diaminetetraacetic acid, 0.02 mg/ml NaN<sub>3</sub>, 8% w/v glycerol,

150  $\mu$ M each of dNTP, 0.25  $\mu$ M each primer, 5U rTth polymerase, 3.25 mM Mn(OAc)<sub>2</sub>, and 5  $\mu$ l blood equivalents of target (see Example 3). Since RNA and the rTth polymerase enzyme are unstable in the presence of Mn(OAc)<sub>2</sub>, the

5 Mn(OAc)<sub>2</sub> should be added just before target addition.

Optimal conditions for cDNA synthesis and thermal cycling readily can be determined by those skilled in the art. The reaction is incubated in a Perkin-Elmer Thermal Cycler 480. Optimal conditions for cDNA synthesis and thermal cycling

10 can readily be determined by those skilled in the art.

Conditions which may be found useful include cDNA synthesis at 60°-70° for 15-45 min, and 30-45 amplification cycles at 94°C, 1 min; 55°C-70°C, 1 min; 72°C, 2 min. One step RT-PCR also may be performed by using a dual enzyme procedure with  
15 Taq polymerase and a reverse transcriptase enzyme, such as MMLV or AMV RT.

B. Traditional RT-PCR. Alternatively, a traditional two step RT-PCR reaction may be performed, as described by K.-Q. Hu et al., Virology 181:721-726 (1991), as follows:

20 The extracted mRNA is transcribed in a 25  $\mu$ l reaction mixture containing 10 mM Tris-HCl, pH 8.3, 5 mM MgCl<sub>2</sub>, 500  $\mu$ M dNTP, 20 U RNasin, 1  $\mu$ M antisense primer, and 25 U AMV (avian myeloblastosis virus) or MMLV (Moloney murine leukemia virus) reverse transcriptase. Reverse  
25 transcription is performed at 37-45°C for 30-60 min, followed by further incubation at 95°C for 5 min to inactivate the RT. PCR is performed using 10  $\mu$ l of the cDNA reaction in a final PCR reaction volume of 50  $\mu$ l containing 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 2 mM MgCl<sub>2</sub>, 200  $\mu$ M dNTP,  
30 0.5  $\mu$ M of each primer and 2.5 U of Taq polymerase. Optimal conditions for cDNA synthesis and thermal cycling can be readily determined by those skilled in the art. The reaction is incubated in a Perkin-Elmer Thermal Cycler 480. Conditions which may be found useful include 30-45 cycles of  
35 amplification (94°C, 1 min; 55°-70°C, 1 min; 72°C, 2 min), final extension (72°C, 10 min) and soak at 4°C.



C. PCR Fragment Analysis. The correct products can then be verified by size determination using gel electrophoresis with fluorescent intercalators or by southern blotting techniques using labeled probes against the internal sequences of the PCR product. The probes may also be polynucleotides analogs, such as morpholinos or peptide nucleic acids (PNA). Detection of a sequence selected from the group consisting of SEQUENCE ID NO 1, or fragments or complements thereof is then indicative of the presence of the TREPA gene, suggesting the diagnosis of inflammatory disease.

D. Automated Quantitative PCR. Quantities of specific PCR products in a given sample can be automatically verified by using a fluorescently labeled probe against the internal sequences of the PCR product coupled to a quenching moiety. Uncoupling of the fluorescent tag to the quenching moiety from the internal probe hybridized to the PCR product by the PCR polymerase during thermocycling can be detected by machines such as the Perkin-Elmer 7700 quantitative PCR machine, and is proportional to the amount of PCR product present in the sample. Detection of a sequence selected from the group consisting of SEQUENCE ID NO 1, or fragments or complements thereof is then indicative of the presence of the TREPA gene, suggesting the diagnosis of inflammatory disease.

#### Example 9: OH-PCR

A. Probe selection and Labeling. Target-specific primers and probes are designed to detect the above target sequence by oligonucleotide hybridization PCR. Publications WO 92/10505, published 25 June 1992 and WO 92/11388 published 9 July 1992 teach methods for labeling oligonucleotides at their 5' and 3' ends, respectively. According to one known method for labeling an oligonucleotide, a label-phosphoramidite reagent is prepared and used to add the label to the oligonucleotide during its synthesis. For example, see N. T. Thuong et al., Tet. Letters 29(46):5905-5908 (1988); or J. S. Cohen et al.,

published U.S. Patent Application 07/246,688 (NTIS ORDER No. PAT-APPL-7-246,688) (1989). Preferably, probes are labeled at their 3' end to prevent participation in PCR and the formation of undesired extension products. For one step OH-PCR, the probe should have a  $T_m$  at least 15°C below the  $T_m$  of the primers. The primers and probes are labeled with either capturable or detectable moieties using standard phosphoramidite chemistry which is well-known to one skilled in the art.

10        B. One Step Oligo Hybridization PCR. OH-PCR is performed on a 200  $\mu$ l reaction containing 50 mM (N,N,-bis[2-Hydroxyethyl]glycine), pH 8.15, 81.7 mM KOAc, 33.33 mM KOH, 0.01 mg/ml bovine serum albumin, 0.1 mM ethylene diaminetetraacetic acid, 0.02 mg/ml NaN<sub>3</sub>, 8% w/v glycerol, 15        150  $\mu$ M each of dNTP, 0.25  $\mu$ M each primer, 3.75 nM probe, 5U rTth polymerase, 3.25 mM Mn(OAc)<sub>2</sub>, and 5  $\mu$ l blood equivalents of target (see Example 3). Since RNA and the rTth polymerase enzyme are unstable in the presence of Mn(OAc)<sub>2</sub>, the Mn(OAc)<sub>2</sub> should be added just before target 20        addition. The reaction is incubated in a Perkin-Elmer Thermal Cycler 480. Optimal conditions for cDNA synthesis and thermal cycling can be readily determined by those skilled in the art. Conditions which may be found useful include cDNA synthesis (60°C, 30 min), 30-45 amplification 25        cycles (94°C, 40 sec; 55-70°C, 60 sec), oligo-hybridization (97°C, 5 min; 15°C, 5 min; 15°C soak). The correct reaction product contains at least one of the strands of the PCR product and an internally hybridized probe.

30        C. OH-PCR product analysis. Amplified reaction products are detected on an LCx® analyzer system (available from Abbott Laboratories, Abbott Park, IL). Briefly, the correct reaction product is captured by an antibody labeled microparticle at a capturable site on either the PCR product strand or the hybridization probe, and the complex is 35        detected by binding of a detectable antibody conjugate to either a detectable site on the probe or the PCR strand. Only a complex containing a PCR strand hybridized with the

internal probe is detectable. The detection of this complex is then indicative of the presence of the TREPA gene, suggesting the diagnosis of inflammatory disease, such as rheumatoid arthritis.

5 Many other detection formats exist which can be used to detect the presence of the TREPA encoding nucleic acid sequence. The sequence may also be detected by other methods including but not limited to ligase chain reaction (LCR, Abbott Laboratories, Abbott Park, IL), Q-beta  
10 replicase (Gene-Trak<sup>TM</sup>, Naperville, Illinois), branched chain reaction (Chiron, Emeryville, CA), and strand displacement assays (Becton Dickinson, Research Triangle Park, NC).

#### 15 Example 10: Synthetic Peptide Production

Synthetic peptides (SEQUENCE ID NOS 10-36 or fragments thereof) are prepared based upon the predicted amino acid sequence of the TREPA polypeptide (see Example 1). All peptides are synthesized on an ABI Peptide Synthesizer  
20 (available from Applied Biosciences, LOCATION), Model 431A, using Fmoc chemistry, standard cycles and DCC-HOBT activation. Cleavage and deprotection conditions are as follows: The resin is added to 20 ml trifluoroacetic acid (TFA), 0.3 ml water, 0.2 ml ethanedithiol, 0.2 ml  
25 thioanisole and 100 mg phenol, and stirred at room temperature for 1.5 hours. The resin then is filtered by suction and the peptide is obtained by precipitation of the TFA solution with ether followed by filtration. Each peptide is purified via reverse-phase preparative HPLC using  
30 a water/acetonitrile/0.1% TFA gradient and lyophilized. The product is confirmed by mass spectrometry (see Example 12).

Disulfide bond formation is accomplished using auto-oxidation conditions, as follows: The peptide is dissolved in a minimum amount of DMSO (approximately 10 ml) before  
35 adding buffer (0.1 M Tris-HCl, pH 6.2) to a concentration of 0.3 - 0.8 mg/ml. The reaction is monitored by HPLC until complete formation of the disulfide bond, followed by reverse-phase preparative HPLC using a water / acetonitrile

/ 0.1% TFA gradient and lyophilization. The product then is confirmed by mass spectrometry (see Example 12).

The purified peptides can be conjugated to Keyhole Limpet Hemocyanin or other immunoreactive molecules with  
 5 glutaraldehyde, mixed with adjuvant, and injected into animals.

Example 11: Expression of Protein in a Cell Line

A. Construction of EST Expression Plasmid. Plasmid  
 10 577, described in U.S. patent application Serial No. 08/478,073, filed June 7, 1995 and incorporated herein by reference, has been constructed for the expression of secreted antigens in a permanent cell line. This plasmid contains the following DNA segments: (a) a 2.3 Kb fragment  
 15 of pBR322 containing bacterial beta-lactamase and origin of DNA replication; (b) a 1.8 Kb cassette directing expression of a neomycin resistance gene under control of HSV-1 thymidine kinase promoter and poly-A addition signals; (c) a  
 20 1.9 Kb cassette directing expression of a dihydrofolate reductase gene under the control of a SV-40 promoter and poly-A addition signals; (d) a 3.5 Kb cassette directing expression of a rabbit immunoglobulin heavy chain signal sequence fused to a modified hepatitis C virus (HCV) E2  
 25 protein under the control of the Simian Virus 40 T-Ag promoter and transcription enhancer, the hepatitis B virus surface antigen (HBsAg) enhancer I followed by a fragment of Herpes Simplex Virus-1 (HSV-1) genome providing poly-A addition signals; and (e) a residual 0.7 Kb fragment of  
 30 Simian Virus 40 genome late region of no function in this plasmid. All of the segments of the vector were assembled by standard methods known to those skilled in the art of molecular biology.

Plasmids for the expression of secretable TREPA  
 35 proteins are constructed by replacing the hepatitis C virus E2 protein coding sequence in plasmid 577 with those from the EST sequence selected from the group consisting of SEQUENCE ID NO 1 or fragments thereof, as follows:  
 Digestion of plasmid 577 with XbaI releases the hepatitis C

EST-23-202-226

virus E2 gene fragment. The resulting plasmid backbone allows insertion of the TREPA insert downstream of the rabbit immunoglobulin heavy chain signal sequence which directs the expressed proteins into the secretory pathway of the cell. The TREPA fragment is generated by PCR using standard procedures. Encoded in the sense PCR primer sequence is an Xba 1 site, immediately followed by a 12 nucleotide sequence that encodes the amino acid sequence Ser-Asn-Glu-Leu ("SNEL") to promote signal protease processing, efficient secretion and final product stability in culture fluids. Immediately following this 12 nucleotide sequence, the primer contains nucleotides complementary to template sequences encoding amino acids of the TREPA sequence. The antisense primer incorporates a sequence encoding the eight amino acids Asp-Tyr-Lys-Asp-Asp-Asp-Lys just before the stop codons. Within this sequence is the recognition site for a monoclonal antibody (MAb) designated anti-FLAG M2 (Eastman Kodak, Co., New Haven, CT). It is incorporated to aid in analysis and purification of the EST protein product. PCR is performed using GeneAmp<sup>®</sup> reagents obtained from Perkin-Elmer-Cetus, essentially as directed by the supplier's instructions. PCR primers are used at a final concentration of 0.5  $\mu$ M. PCR is performed on the pSPORT1 plasmid template in a 100  $\mu$ l reaction for 35 cycles (94°C, 30 seconds; 55°C, 30 seconds; 72°C, 90 seconds) followed by an extension cycle of 72°C for 10 min.

Another example of a plasmid for the expression of secretable TREPA proteins is created by fusing in-frame sequence encoding a modified cytomegalovirus UL4 gene leader sequence (SEQUENCE ID NO 4) to the synthetic octapeptide encoding the eight amino acids eight amino acids Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (recognized by the monoclonal antibody designated anti-FLAG M2; SEQUENCE ID NO 5) followed by an appropriate carboxyl-terminal portion of TREPA which interacts with the cognate receptor, and can therefore induce biological activity (SEQUENCE ID NO 3). One such form of soluble TRAIL is presented in SEQUENCE ID NO 3.

This engineered synthetic open reading frame (SEQUENCE ID NO 4 + SEQUENCE ID NO 5 + SEQUENCE ID NO 3) can be placed in an appropriate DNA backbone as is appropriate for the intended target cell, such as pcDNA3 (Invitrogen Corp, San Diego, CA).

B. Transfection of Dihydrofolate Reductase Deficient Chinese Hamster Ovary Cells. The plasmid described supra is transfected into CHO/dhfr- cells (DXB-111) (Uriacio, et al., PNAS 77, 4451-4466 ; 1980); these cells are available from the A.T.C.C., 12301 Parklawn Drive, Rockville, MD 20852, under Accession No. CRL 9096), using the cationic liposome-mediated procedure (P. L. Felgner et al., PNAS 84:7413-7417 (1987), as follows: CHO/dhfr- cells are cultured in Ham's F-12 media supplemented with 10% fetal calf serum, L-glutamine (1 mM) and freshly seeded into a 25 cm<sup>2</sup> flask at a density of 5 - 8 x 10<sup>5</sup> cells per flask. The cells are grown to between 60 and 80% confluency for transfection. Twenty micrograms of plasmid DNA are added to 1.5 ml of Opti-MEM I medium and 100 µl of Lipofectin Reagent (Gibco-BRL; Grand Island, NY) are added to a second 1.5 ml portion of Opti-MEM I media. The two solutions are mixed and incubated at room temperature for 20 min. The culture medium is removed from cells, and the cells are rinsed 3 times with 5 ml of Opti-MEM I medium. The Opti-MEM I-Lipofection-plasmid DNA solution is then overlaid onto the cells. The cells are incubated for 3 h at 37°C, after which time the Opti-MEM I-Lipofectin-DNA solution is replaced with culture medium for an additional 24 h prior to selection.

C. Selection and Amplification. One day after transfection, cells are passaged 1:3 and incubated with dhfr/G418 selection medium (hereafter, "F-12 minus medium G"). Selection medium is Ham's F-12 with L-glutamine and without hypoxanthine, thymidine, and glycine (JRH Biosciences, Lenexa, Kansas), and 300 µg per ml G418 (Gibco-BRL; Grand Island, NY). Media volume to surface area ratios of 5 ml per 25 cm<sup>2</sup> are maintained. After approximately two

weeks, DHFR/G418 cells are expanded to allow passage and continuous maintenance in F-12 minus medium G.

Amplification of each of the transfected EST genes is achieved by stepwise selection of DHFR<sup>+</sup>, G418<sup>+</sup> cells with  
5 methotrexate (reviewed by R. Schimke, Cell 37:705-713 [1984]). Cells are incubated with F-12 minus medium G containing 150 nM methotrexate (MTX) (Sigma, St. Louis, MO) for approximately two weeks until resistant colonies appear. Further gene amplification is achieved by selection of 150  
10 nM adapted cells with 5  $\mu$ M MTX.

D. Protein/Antigen Production. F-12 minus medium G supplemented with 5  $\mu$ M MTX is overlaid onto just confluent monolayers for 12 to 24 h at 37°C in 5% CO<sub>2</sub>. The growth  
15 medium is removed and the cells are rinsed 3 times with Dulbecco's phosphate buffered saline (PBS) (with calcium and magnesium) (Gibco-BRL; Grand Island, NY), to remove the remaining media/serum which might be present. Cells then are incubated with VAS custom medium (VAS custom formulation with L-glutamine with HEPES without phenol red, available  
20 from JRH Bioscience; Lenexa, KS, product number 52-08678P), for 1 h at 37°C in 5% CO<sub>2</sub>. Cells then are overlaid with VAS for production at 5 ml per T 25 cm<sup>2</sup> flask. The medium is removed after 7 days of incubation and then frozen to await purification with harvests 2, 3 and 4. The monolayers are  
25 overlaid with VAS for 3 more 7-day harvests.

E. Analysis of TREPA Protein/Antigen Expression. Aliquots of VAS supernatants from the cells expressing the engineered protein construct are analyzed either by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) using standard  
30 methods and reagents known in the art (Laemmli discontinuous gels) or by mass spectrometry.

F. Purification. Purification of the EST protein containing the FLAG sequence is performed by immunoaffinity chromatography using an affinity matrix comprising anti-FLAG  
35 M2 monoclonal antibody covalently attached to agarose by hydrazide linkage (Eastman Kodak Co., New Haven, CT). Prior to affinity purification, protein in pooled VAS medium

harvests from roller bottles is exchanged into 50 mM Tris-HCl pH 7.5, 150 mM NaCl buffer using a Sephadex G-25 (Pharmacia Biotech Inc., Uppsala, Sweden) column. Protein in this buffer is applied to the anti-FLAG M2 antibody affinity column, non-binding protein is eluted by washing  
5 the column with 50 mM Tris-HCl pH 7.5, 150 mM NaCl buffer and bound protein is eluted using an excess of FLAG peptide in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl. The excess FLAG peptide can be removed from the EST purified protein by gel  
10 electrophoresis.

Then, the largest cloned insert containing the EST region is sub-cloned into either (i) a eukaryotic expression vector which may contain a cytomegalovirus (CMV) promoter and/or protein fusable sequences which aid in protein expression and detection, or (ii) a bacterial expression vector containing a superoxide-dismutase (SOD) and CMP-KDO synthetase (CKS) or other protein fusion gene for expression of the protein sequence. Methods and vectors which are useful for the production of polypeptides which contain fusion sequences of SOD are described in EPO 0196056, published October 1, 1986, and those of CKS are described in EPO Publication No. 0331961, published September 13, 1989. The purified protein can be used in a variety of techniques, including but not limited to animal immunization studies, solid phase immunoassays, etc.

### Example 12: Chemical Analysis of TREPA Proteins

### A. Analysis of Tryptic Peptide Fragments Using MS.

Serum from a patient with inflammatory disease is run on a polyacrylamide gel using standard procedures and stained with Coomassie Blue. Sections of the gel suspected of containing the unknown polypeptide are excised and subjected to an in-gel reduction, acetamidation, and tryptic digestion. P. Jeno et al, Anal. Bio. 224:451-455 (1995), and J. Rosenfeld et al, Anal. Bio. 203:173-179 (1992). The gel sections are washed with 100 mM  $\text{NH}_4\text{HCO}_3$  and acetonitrile. The shrunken gel pieces are swollen in digestion buffer (50 mM  $\text{NH}_4\text{HCO}_3$ , 5 mM  $\text{CaCl}_2$ , and 12.5  $\mu\text{g/ml}$



trypsin) at 4°C for 45 min. The supernatant is aspirated and replaced with 5 to 10 µl of digestion buffer without trypsin and allowed to incubate overnight at 37°C .

Peptides are extracted with 3 changes of 5% formic acid and acetonitrile, and evaporated to dryness. The peptides are adsorbed to approximately 0.1 µl of POROS R2 sorbent (Perseptive Biosystems, Framingham, Massachusetts) trapped in the tip of a drawn gas chromatography capillary tube by dissolving them in 10 µl of 5% formic acid and passing the solution through the capillary. The adsorbed peptides are washed with water and eluted with 5% formic acid in 60% methanol. The eluant is passed directly into the spraying capillary of an API III mass spectrometer (Perkin-Elmer Sciex, Thornhill, Ontario, Canada) for analysis by nano-electrospray mass spectrometry. M. Wilm et al., Int. J. Mass Spectrom. Ion Process 136:167-180 (1994), and M. Wilm et al., Anal. Chem. 66:1-8 (1994). The masses of the tryptic peptides are determined from the mass spectrum obtained off the first quadrupole. Masses corresponding to predicted peptides can be further analyzed in MS/MS mode to give the amino acid sequence of the peptide.

B. Peptide Fragment Analysis Using LC/MS. The presence of polypeptides predicted from mRNA sequences found in hyperplastic disease tissues also can be confirmed using liquid chromatography/tandem mass spectrometry (LC/MS/MS). D. Hess et al., METHODS, A Companion to Methods in Enzymology 6:227-238 (1994). The serum specimen or tumor extract from the patient is denatured with SDS and reduced with dithiothreitol (1.5 mg/ml) for 30 min at 90°C followed by alkylation with iodoacetamide (4 mg/ml) for 15 min at 25°C. Following acrylamide electrophoresis, the polypeptides are electroblotted to a cationic membrane and stained with Coomassie Blue. Following staining, the membranes are washed and sections thought to contain the unknown polypeptides are cut out and dissected into small pieces. The membranes are placed in 500 µl microcentrifuge tubes and immersed in 10 to 20 µl of proteolytic digestion

buffer (100 mM Tris-HCl, pH 8.2, containing 0.1 M NaCl, 10% acetonitrile, 2 mM CaCl<sub>2</sub>, and 5 ug/ml trypsin) (Sigma, St. Louis, MO). After 15 h at 37°C, 3 µl of saturated urea and 1 µl of 100 µg/ml trypsin are added, and incubated for an additional 5 h at 37°C. The digestion mixture is acidified with 3 µl of 10% trifluoroacetic acid and centrifuged to separate supernatant from membrane. The supernatant is injected directly onto a microbore, reverse phase HPLC column and eluted with a linear gradient of acetonitrile in 0.05% trifluoroacetic acid. The eluate is fed directly into an electrospray mass spectrometer, after passing through a stream splitter if necessary to adjust the volume of material. The data is analyzed following the procedures set forth in Example 12, section A.

15

### Example 13: Gene Immunization Protocol

A. In Vivo Antigen Expression. Gene immunization circumvents protein purification steps by directly expressing an antigen in vivo after inoculation of the appropriate expression vector. Also, production of antigen by this method may allow correct protein folding and glycosylation since the protein is produced in mammalian tissue. The method utilizes insertion of the gene sequence into a plasmid which contains a CMV promoter, expansion and purification of the plasmid, and injection of the plasmid DNA into the muscle tissue of an animal. See, for example, H. Davis et al., Human Molecular Genetics 2:1847-1851 (1993). After one or two booster immunizations, the animal can then be bled, ascites fluid collected or spleen harvested for production of hybridomas.

B. Plasmid Preparation and Purification. EST DNA sequences are generated from the pSPORT1 EST vector using appropriate PCR primers containing suitable 5' restriction sites. The PCR product is cut with appropriate restriction enzymes and inserted into a vector which contains the CMV promoter (for example, pRc/CMV or pcDNA3 vectors from Invitrogen, San Diego, CA). This plasmid then is expanded

35

in the appropriate bacterial strain and purified from the cell lysate using a CsCl gradient or a Qiagen plasmid DNA purification column. All these techniques are familiar to one of ordinary skill in the art of molecular biology.

5        C. Immunization Protocol. Anesthetized animals are immunized intramuscularly with 0.1-100 µg of the purified plasmid diluted in PBS or other DNA uptake enhancers (Cardiotoxin, 25% sucrose). See, for example, H. Davis, et al, Human Gene Therapy 4:733-740 (1993); and P. W. Wolff et al, Biotechniques 11:474-485 (1991). One to two booster  
10 injections are given at monthly intervals.

D. Testing and Use of Antiserum. Animals are bled and the sera tested for antibody using peptides synthesized from the known gene sequence (see Example 16) such as western  
15 blotting or EIA techniques. Antisera produced by this method can then be used to detect the presence of the antigen in a patient's sera or tumor tissue extract by ELISA or Western blotting techniques.

20        Example 14: Production of Antibodies Against TREPA

A. Production of Polyclonal Antisera. Antiserum against TREPA is prepared using peptides whose sequence is derived from that of the TREPA DNA sequence (SEQUENCE ID NO 1). The synthesis of the peptides is described in Example  
25 10.

      1. Peptide Conjugation. Peptide is conjugated to maleimide activated keyhole limpet hemocyanine (KLH, commercially available as Imject®, available from Pierce Chemical Company, Rockford, IL). Imject® contains about 250  
30 moles of reactive maleimide groups per mole of hemocyanine. The activated KLH is dissolved in phosphate buffered saline (PBS, pH 8.4) at a concentration of about 7.7 mg/ml. The peptide is conjugated through the C-terminal cysteine previously added to the synthesized peptide in order to  
35 provide a monospecific point of attachment. The peptide is dissolved in dimethyl sulfoxide (DMSO, Sigma Chemical Company, St. Louis, MO) and reacted with the activated KLH at a mole ratio of about 1.5 moles of peptide per mole of

reactive maleimide attached to the KLH. A procedure for the conjugation of peptides is provided hereinbelow. It is known to the ordinary artisan that the amounts, times and conditions of such a procedure can be varied to optimize peptide conjugation.

The conjugation reaction described hereinbelow is based on obtaining 3 mg of KLH peptide conjugate, which contains about 0.77  $\mu$ moles of reactive maleimide groups. This quantity of peptide conjugate usually is adequate for one primary injection and four booster injections for production of polyclonal antisera in a rabbit. The amount of peptide conjugate produced can be increased or decreased, as appropriate. Briefly, peptide is dissolved in DMSO at a concentration of 1.16  $\mu$ moles/100  $\mu$ l (4.1 mg/100  $\mu$ l) of DMSO. One hundred microliters (100  $\mu$ l) of the DMSO solution is added to 380  $\mu$ l of the activated KLH solution prepared as described hereinabove, and 20  $\mu$ l of PBS (pH 8.4) is added to bring the volume to 500  $\mu$ l. The reaction is incubated overnight at room temperature with stirring. The extent of reaction is determined by measuring the amount of unreacted thiol in the reaction mixture. The difference between the starting concentration of thiol (11.6 mM) and the final concentration is assumed to be the concentration of peptide which has coupled to the activated KLH. The amount of remaining thiol is measured using Ellman's reagent (5,5'-dithiobis(2-nitrobenzoic acid), Pierce Chemical Company, Rockford, IL). Cysteine standards are made at a concentration of 0, 0.1, 0.5, 2, 5, and 20 mM by dissolving 35 mg of cysteine HCl (Pierce Chemical Company, Rockford, IL) in 10 ml of PBS (pH 7.2) and diluting the stock solution to the desired concentration(s). The photometric determination of the concentration of thiol is accomplished by placing 200  $\mu$ l of PBS (pH 8.4) in each well of an Immulon 2  $\otimes$  microwellplate (Dynex Technologies, Chantilly, VA). Next, 10  $\mu$ l of standard or reaction mixture is added to each well. Finally, 20  $\mu$ l of Ellman's reagent at a concentration of 1 mg/ml in PBS (pH 8.4) is added to each well. The wells

are incubated for 10 minutes at room temperature, and the absorbance of all wells is read at 415 nm with a microplate reader (such as the BioRad Model 3550, BioRad, Richmond, CA). The absorbance of the standards is used to construct a standard curve and the thiol concentration of the reaction mixture is determined from the standard curve. A decrease in the concentration of free thiol is indicative of a successful conjugation reaction. Unreacted peptide is removed by dialysis against PBS (pH 7.2) at room temperature for 6 hours. The conjugate is stored at 2-8°C if it is to be used immediately; otherwise, it is stored at -20°C or colder.

2. Animal Immunization. Female white New Zealand rabbits weighing 2 kg or more are used for raising polyclonal antiserum. Generally, one animal is immunized per peptide conjugate prepared as described hereinabove. One week prior to the first immunization, 5 to 10 ml of blood is obtained from the animal to serve as a non-immune prebleed sample. The primary immunogen utilized for the primary immunization consists of 1 mg of peptide conjugate prepared as described hereinabove emulsified in complete Freund's adjuvant (CFA) (Difco, Detroit, MI). The primary immunogen is prepared by taking 0.5 ml of the peptide conjugate diluted to a concentration of 2 mg/ml and emulsifying it with 0.5 ml of CFA. The immunogen is injected into several sites of the animal, and injections can include subcutaneous, intraperitoneal and intramuscular. Four weeks following the primary immunization, a booster immunization is administered utilizing a second immunogen consisting of 0.5 mg of peptide conjugate emulsified in incomplete Freund's adjuvant (IFA, available from Difco, Detroit, MI). The immunogen used for the booster immunization dose is prepared by emulsifying 0.5 ml of peptide conjugate diluted to 1 mg/ml with 0.5 ml of IFA. Again, the booster dose is administered into several sites and can utilize subcutaneous, intraperitoneal and intramuscular types of injections. The animal is bled (5

ml) two weeks after the booster immunization, and the serum is tested for immunoreactivity to the peptide, as described below. The booster and bleed schedule is repeated at 4 week intervals until an adequate titer is obtained. The titer or  
5 concentration of antiserum is determined by microtiter EIA as described in Example 17, below. An antibody titer of 1:500 or greater is considered an adequate titer for further use and study.

B. Production of Monoclonal Antibody.

10 1. Immunization Protocol. Mice are immunized using the immunogen prepared as described hereinabove, except that the amount of the peptide conjugate for monoclonal antibody production in mice is one-tenth the  
15 amount of the peptide conjugate used to produce polyclonal antisera in rabbits. Thus, the primary immunogen consists of 100 µg of peptide conjugate in 0.1 ml of CFA, while the immunogen used for booster immunizations consists of 50 µg of peptide conjugate in 0.1 ml of IFA. Hybridomas for the generation of monoclonal antibodies are prepared and  
20 screened using standard techniques. The methods used for monoclonal antibody development follow procedures known in the art and detailed in Kohler and Milstein, Nature 256:494 (1975), and reviewed in J.G.R. Hurrel, ed., Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC  
25 Press, Inc., Boca Raton, FL (1982). Another method of monoclonal antibody development which is based on the Kohler and Milstein method is that of L.T. Mimms et al., Virology 176:604-619 (1990), which is incorporated herein by reference.

30 The immunization regimen (5 mice) consists of a primary immunization with additional booster immunizations. The primary immunogen used for the primary immunization consists of 500 µg of peptide conjugate in 250 µl of PBS (pH 7.2) emulsified with 250 µl of CFA. Booster immunizations  
35 performed at approximately two weeks and four weeks post primary immunization consist of 250 µg of peptide conjugate in 250 µl of PBS (pH 7.2) emulsified with 250 µl IFA.

Immunogen used for booster immunizations is inoculated intraperitoneally and subcutaneously into each mouse, with each mouse receiving a total of 100  $\mu$ l of this so-prepared immunogen per immunization. Individual mice are screened  
5 for immune response by microtiter plate enzyme immunoassay (EIA) as described in Example 17 approximately four weeks after the third immunization. Mice are inoculated intravenously with 50  $\mu$ g of peptide conjugate in PBS (pH 7.2) approximately fifteen weeks after the third  
10 immunization.

Three days after this intravenous boost, splenocytes are fused with, for example, Sp2/0-Ag14 myeloma cells from Milstein Laboratories, England, using the polyethylene glycol (PEG) method. The fusions are cultured in Iscove's  
15 Modified Dulbecco's Medium (IMDM) containing 10% fetal calf serum (FCS), plus 1% hypoxanthine, aminopterin and thymidine (HAT). Bulk cultures are screened by microtiter plate EIA following the protocol in Example 17. Clones reactive with the peptide used as the immunogen and non-reactive with  
20 other peptides are selected for final expansion. Clones thus selected are expanded, aliquoted and frozen in IMDM containing 10% FCS and 10% dimethylsulfoxide.

2. Production of Ascites Fluid Containing Monoclonal Antibodies. Frozen hybridoma cells prepared as described  
25 hereinabove are thawed and placed into expansion culture. Viable hybridoma cells are inoculated intraperitoneally into Pristane treated mice. Ascitic fluid is removed from the mice, pooled, filtered through a 0.2  $\mu$  filter and subjected to an immunoglobulin class G (IgG) analysis to determine the  
30 volume of the Protein A column required for the purification.

3. Purification of Monoclonal Antibodies From Ascites Fluid. Briefly, filtered and thawed ascites fluid is mixed with an equal volume of Protein A sepharose binding buffer  
35 (1.5M glycine, 3.0M NaCl, pH 8.9) and refiltered through a 0.2  $\mu$  filter. The volume of the Protein A column is determined by the quantity of IgG present in the ascites

fluid. The eluate then is dialyzed against PBS pH 7.2 overnight at 2-8°C. The dialyzed monoclonal antibody is sterile filtered and dispensed in aliquots. The immunoreactivity of the purified monoclonal antibody is confirmed by determining its ability to specifically bind to the peptide (SEQUENCE ID NO 6) used as the immunogen by use of the EIA microtiter plate assay procedure of Example 17. The specificity of the purified monoclonal antibody is confirmed by determining its lack of binding to irrelevant peptides. The purified anti-TREPA monoclonal thus prepared and characterized is placed at either 2-8°C for short term storage and at -80°C for long term storage.

4. Further Characterization of Monoclonal Antibody. The isotype and subtype of the monoclonal antibody produced as described hereinabove can be determined using commercially available kits (available from Amersham, Inc., Arlington Heights, IL). Stability testing also can be performed on the monoclonal antibody by placing an aliquot of the monoclonal antibody in continuous storage at 2-8°C and assaying optical density (OD) readings throughout the course of a given period of time.

C. Use of Recombinant Proteins as Immunogens. It is within the scope of the present invention that recombinant proteins made as described herein can be utilized as immunogens in the production of polyclonal and monoclonal antibodies, with corresponding changes in reagents and techniques known to those skilled in the art.

D. Immunogen Variation. It is contemplated that peptide not conjugated to a carrier such as described hereinabove in 1(A) may be utilized at times as the immunogen for animal inoculation. This so-prepared immunogen may be attached to adjuvants such as complete or incomplete Freund's adjuvant.

35 Example 15: Purification of TREPA peptide specific antibodies from serum

Immune sera is affinity purified using immobilized synthetic peptides by methods known in the art. Antiserum



produced against a peptide as described in Example 10 is affinity purified in a variety of ways. An IgG fraction is obtained by passing the diluted, crude antiserum over a Protein A column (Affi-Gel protein A, Bio-Rad, Hercules, CA). Elution with Binding Buffer supplied by the manufacturer removes all proteins that are not immunoglobulins. Elution with pH 3, buffered glycine, 0.1M, gives an immunoglobulin preparation that is substantially free of albumin and other serum proteins.

Immunoaffinity chromatography is performed to obtain a preparation with a higher fraction of specific antigen-binding antibody. The peptide used to raise the antiserum is immobilized on a chromatography resin and the specific antibodies directed against its epitopes are adsorbed to the resin. After washing away non-binding components, the specific antibodies are eluted with 0.1 M glycine buffer, pH 2.3 antibody fractions are immediately neutralized with 1.0M Tris buffer, pH 8.0, to preserve immunoreactivity. The resin chosen depends on the reactive groups present in the peptide. If the peptide has an amino group, a resin such as Affi-Gel 10 or Affi-Gel 15 is used (Bio-Rad, Hercules, CA). If coupling through a carboxy group on the peptide is desired, Affi-Gel 102 can be used (Bio-Rad, Hercules, CA). If the peptide has a free sulfhydryl group, an organomercurial resin such as Affi-Gel 501 can be used (Bio-Rad, Hercules, CA).

Alternatively, spleens can be harvested and used in the production of hybridomas to produce monoclonal antibodies.

#### 30      Example 16: Western Blotting of tissue samples

Tissue samples are homogenized in SDS-PAGE sample buffer (50 mM Tris-HCl, pH 6.8, 100 mM dithiothreitol, 2% SDS, 0.1% bromophenol blue, 10% glycerol), heated at 100°C for 10 min and run on a 14% SDS-PAGE with a 25 mM Tris-HCl, pH 8.3, 250 mM Glycine, 0.1% SDS running buffer. The proteins are electrophoretically transferred to nitrocellulose in a transfer buffer containing 39 mM glycine, 48 mM Tris-HCl, pH 8.3, 0.037% SDS, 20% methanol.

The nitrocellulose is dried at room temperature for 60 min and then blocked with a PBS solution containing either bovine serum albumin or 5% nonfat dried milk for 2 h at 4°C.

The filter is placed in a heat-sealable plastic bag  
5 containing a solution of 5% nonfat dried milk in PBS with a 1:100 to 1:2000 dilution of affinity purified anti-EST peptide antibodies (see Example 15), incubated at 4°C for 2 h, followed by 3 10 min washes in PBS. An alkaline phosphatase conjugated secondary antibody (i.e., anti-  
10 mouse/rabbit IgG), is added at a 1:200 to 1:2000 dilution to the filter in a 150 mM NaCl, 50 mM Tris-HCl, pH 7.5 buffer and incubated for 1h at room temperature.

The bands are visualized upon the addition and development of a chromogenic substrate such as 5-bromo-4-  
15 chloro-3-indolyl phosphate/nitro blue tetrazolium (BCIP/NBT). This chromogenic solution contains 0.033% NBT and 0.016% BCIP in a solution containing 100 mM NaCl, 5 mM MgCl<sub>2</sub> and 100 mM Tris-HCl, pH 9.5. The filter is incubated in the solution at room temperature until the bands develop  
20 to the desired intensity. Development is stopped in a PBS buffer containing 2 mM EDTA. Molecular mass determination is made based upon the mobility of pre-stained molecular weight standards (Rainbow markers, Amersham, Arlington Heights, IL).

25

Example 17: EIA Microtiter Plate Assay

To demonstrate how relative immunoreactivity for EST synthetic peptides is measured, wells of 96-well microtiter plates (Dynatec Immunolon 4 polystyrene) are coated for 16 h at 4°C with 100 µl of the EST peptide at the following concentrations: 500 µM, 50 µM, 5 µM, 0.5 µM, 0.05 µM, and 0.005 µM. The buffer used for the application of these peptides is 100 mM morpholino-ethane sulfonic acid, pH 5.5. The EST peptides coated wells are then washed 3 times with wash buffer (8 mM sodium phosphate, 2 mM potassium phosphate, 140 mM sodium chloride, 10 mM potassium chloride, 0.05% Tween 20, 0.1% bovine serum albumin, pH 7.4).

The wells then are blocked for 1 h at room temperature with 9% w/v of Carnation skim milk powder in phosphate buffered saline (8 mM sodium phosphate, 2 mM potassium phosphate, 140 mM sodium chloride, 10 mM potassium chloride, pH 7.4). The wells next are washed 3 times with wash buffer.

The test specimen or control (mouse or rabbit antiserum) is diluted 150-fold with 4.5% Carnation skim milk powder in PBS. Then, 100 µl of this sample are incubated in the wells at 37°C for 1 h, followed by 3 washes with wash buffer.

Horseradish peroxidase conjugated goat anti-mouse/rabbit IgG is used as a second antibody label to bind with the anti-EST-antibody/EST antigen complex formed in positive wells. 100 µL of HRPO-goat anti-mouse/rabbit IgG conjugate at a dilution of about 1:5000 in wash buffer are added to each well and incubated at room temperature for 1h. The wells are washed 3 times with wash buffer.

Positive wells are identified by the absorbance readings at 405 nm after exposure of the wells to 100 µL of ABTS solution (2,2'-azinobis-[3-ethylbenzothiazoline-6-sulfonic acid] diammonium salt) (Pierce Chemical Co., Rockford, IL, USA). Alternatively, color development can be achieved with the addition to each well of 100 µl of a solution of o-phenylene diamine (OPD) in hydrogen peroxide

and a 10 min incubation at room temperature. The color development reaction is quenched with 100 µl of 1N sulfuric acid. The colors in the wells are read as absorbance with a Dynatech MR5000 plate reader at 490 nm and 630 nm

5 wavelengths. A positive signal is indicative the presence of anti-EST peptide antibodies.

#### Example 18: Coating of Solid Phase Particles

##### A. Coating of Microparticles with Anti-EST

10 Peptide Antibody. Affinity purified anti-EST peptide antibodies (see Example 15) are coated onto microparticles which may include polystyrene, carboxylated polystyrene, polymethylacrylate or similar particles with a radius in the range of about 0.1 to 20 µm. Microparticles may be either  
15 passively or actively coated. One method is coating of EDAC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Aldrich Chemical .Co., Milwaukee, WI) activated carboxylated latex microparticles with anti-EST antibody. Briefly, a final 0.375% solid solution of resin washed  
20 carboxylated latex microparticles are mixed in a solution containing 50 mM MES buffer, pH 4.0 and 150 mg/l of affinity purified anti-EST antibody (see Example 15) for 15 min in an appropriate container. EDAC coupling agent is added to a final concentration of 5.5 µg/ml to the mixture and mixed  
25 for 2.5 h at room temperature.

The microparticles then are washed with 8 volumes of a Tween 20®/sodium phosphate, pH 7.2 wash buffer by tangential flow filtration using a 0.2 µm Microgon Filtration module. Washed microparticles are stored in an appropriate buffer,  
30 usually containing a dilute surfactant and irrelevant protein as a blocking agent, until needed.

B. Coating of 1/4 inch Beads. Anti-EST antibodies also may be coated on the surface of 1/4 inch polystyrene beads by routine methods known in the art (Snitman et al, US  
35 Patent 5,273,882, incorporated herein by reference) and used in competitive binding or EIA sandwich assays. Polystyrene beads are first cleaned by ultrasonically cleaning them for about 15 seconds in 10 mM NaHCO<sub>3</sub> buffer at pH 8.0. The beads are

then washed in deionized water until all fines are removed. Beads are then immersed in an antibody solution in 10 mM carbonate buffer, pH 8 to 9.5. The antibody solution can be as dilute as 1 µg/ml in the case of high affinity monoclonal antibodies or as concentrated as about 500 µg/ml for polyclonal antibodies which have not been affinity purified. Beads are coated for at least 12 hours at room temperature and then washed with deionized water. Beads may be dried or stored wet. They may be overcoated with protein stabilizers (sucrose) or non-specific binding blockers (irrelevant proteins, Carnation skim milk, or the like).

Example 19: Microparticle Enzyme Immunoassay (MEIA)

EST proteins and peptides are detected using a standard commercialized antigen competition EIA assay or polyclonal antibody sandwich EIA assay on the IMx® Analyzer (Abbott Laboratories, Abbott Park, IL). Briefly, samples suspected of containing the EST protein are incubated in the presence of anti-EST coated microparticles (see Example 16). The microparticles are washed and secondary polyclonal anti-EST antibodies conjugated with detectable entities (i.e., alkaline phosphatase) are added and incubated with the microparticles. The microparticles are washed and the bound antibody/antigen/antibody complexes are detected by adding a substrate (i.e. 4-methyl umbelliferyl phosphate) (MUP) that will react with the secondary conjugated antibody to generate a detectable signal. An elevated signal, indicating the presence of EST protein, is diagnostic of cancer.

A competitive binding assay uses a detectably labeled peptide that generates a specific background signal on the IMx® analyzer when the peptide is bound to an anti-peptide antibody coated microparticle. The labeled peptide also is added to the microparticles in the presence of patient samples suspected of containing the EST protein. The EST protein in the patient sample will compete with the labeled EST peptide for binding sites on the microparticle resulting in lowered IMx® signals. A lowered signal, indicating the

The TREPA gene polynucleotides and the proteins encoded therein which are provided and discussed hereinabove therefore are thought to be useful as markers of TREPA-associated disease. Tests based upon the appearance of this marker in a test sample such as blood, plasma or serum can provide low cost, non-invasive, diagnostic information to aid the physician to make a diagnosis of cancer, to help select a therapy protocol, or to monitor the success of the chosen therapy. This marker may appear in readily accessible body fluids such as blood, urine or stool as antigens derived from the diseased tissue which are detectable by immunological methods. This marker may be elevated in a disease state, altered in a disease state, or be a normal protein which appears in an inappropriate body compartment. Also, the TREPA gene polynucleotides and the proteins encoded thereby, provided by the present invention, are useful for treating a variety of disorders.

Example 20: HUVEC Experiment

Ligand preparation: UL4flagTREPA was transiently transfected into COS-7 cells using lipafectamine. Three days after the media was changed, 2 ml of supernatant were collected from the cells. The supernatant was centrifuged at 1000rpm to remove cell debris, and the protein was concentrated by centrifugation on a centricon 10 column (Pierce Chemical, Rockford, IL) at 5000 X g for 1 hour. As a negative control, supernatants from cells treated the same way, except for the absence of DNA, were similarly prepared (moc).

Plate preparation: Wells of a 96 well tissue culture plate were coated with 10ug/ml of m2 anti flag antibody (Kodak) in 100ul PBS for 3 hours. The wells were washed three times with PBS and coated with the following overnight: Two wells were coated with 100ul concentrated supernatant from moc transfected cells. All wells were washed with PBS three times before the addition of HUVEC (human umbilical endothelium cell) cells.

Cells: One 10 cm plate of 80% confluent passage 4 HUVEC cells was washed with PBS and then incubated with 5ml PBS for 3 minutes at 37C to remove the cells. 10 ml of serum free media was added, and the cells were counted and centrifuged at 1500 X g. Cells were resuspended in 10ml of serum free media and counted. These cells were seeded at 1500 cells per well in 100 ul EGM media (Clonetics, Walkersville, MD) with 2% fetal bovine serum. The plate was incubated at 3% CO<sub>2</sub>, 37C for six days.

Results: The wells were washed with PBS and stained with crystal violet stain (20% methanol, 0.1% crystal violet). The wells were washed five times with water and photographed on a microscope with 10 fold magnification. The number of cells per photograph were counted. The results are as follows:

moc	UL4flagTREPA
114	861
103	661

5           In another experiment, HUVEC cells were similarly treated except that they were cultured in EGM media without EGF and hydrocortisone, and 1% fetal bovine serum. Their numbers were measured by staining with the fluorescent dye calcien-AM (Molecular Probes, Eugene, OR). Fluorescence was measured with an excitation wavelength of 485 nm and read at 530 nm on a cytofluor 2300 system at sensitivity 2 (Millipore, Bedford, MA). Background fluorescence from wells with media and no cells were subtracted, and the results are as follows:

15	moc	UL4flag TREPA
	-1	41
	0	38
	-1	41
20	-1	41

These results indicate that UL4flagTREPA is capable of inducing proliferation in HUVEC cells as measured both by direct cell counting and indirect fluorescent staining.

#### Example 21: Proliferation of Three Additional Types of Cells

Ligand preparation: A soluble construct of UL4flagTREPA (see SEQUENCE ID NO 3) was transfected into CHO cells.

Clones were selected using geneticin selection at 500 ug/ml and assayed for the production of soluble flagTREPA by

western blot of the conditioned media. One flagTREPA expressing clone was propagated and used to produce conditioned media. 500 ml of conditioned media was collected and centrifuged at 1000 rpm to remove cell debris. Two ml of M2 anti-flag conjugated sepharose were added to the supernatant and incubated with gentle agitation over night at 4 degrees C. The media was centrifuged at 1000 rpm to



harvest the beads. The media was discarded. The beads were washed with 40 ml of PBS containing 2 mM  $MgCl_2$ . 1mM flag peptide was prepared in PBS with 2 mM  $MgCl_2$  and incubated with the beads for ten minutes. Seven 0.5 ml fractions were eluted and analyzed by SDS-PAGE. The first four fractions were pooled and dialyzed for 4 hours in 500 ml PBS with 2 mM  $MgCl_2$  with one change of buffer after the first two hours to remove the flag peptide. The resulting protein was analyzed by SDS PAGE and the protein was quantitated by spectrophotometer using the GeneQuant DNA/RNA calculator (Pharmacia Biotech) total protein function.

Cells: Three additional endothelial cell types were assayed, HA (human aortic endothelial cells), HBE (Human Brain microvascular endothelial cells) and HUMVE (Human microvascular Endothelial cells). All three cells types were similarly treated. One 10 cm plate of 80% confluent passage under 9 was washed in PBS and incubated with 2 ml PBS for 3 minutes at 37C to remove cells. The cells were washed in 10 ml serum free media and counted. Cells were seeded at 1500 cells per well in 100 ul EGM media without EGF and Hydrocortisone, and 1% fetal bovine serum. 5ng (final concentration of 50 ng/ml) of purified TREPA was added to each well. Five days later, cell numbers were measured by staining with the fluorescent dye calcien-AM (molecular Probes, Eugene, Or.). Fluorescence was measured with an excitation wavelength of 485 nm and read at 530 nm on a cytofluor 2300 system at sensitivity 2 (Millipore, Bedford, MA). Background fluorescence from wells with media and no cells were subtracted, and the results are as follows:

untreated HUMVE cells	UL4flagTREPA treated HUMVE
6	26
6	25
5	21
5	21

The above results indicate that purified UL4flagTREPA is capable of inducing proliferation in a variety of endothelial cells as measured by indirect fluorescent staining. These cells include both large vessel endothelium (HA) and microvasculature endothelium (HBA and HUMVE).

The above results indicate that purified UL4flagTREPA is capable of inducing proliferation in a variety of endothelial cells as measured by indirect fluorescent staining. These cells include both large vessel endothelium (HA) and microvasculature endothelium (HBA and HUMVE).

## SEQUENCE LISTING

## (1) GENERAL INFORMATION

- (i) APPLICANT: Wiley, Steven R.
- (ii) TITLE OF THE INVENTION: MEMBER OF THE TNF  
FAMILY USEFUL FOR TREATMENT AND DIAGNOSIS OF  
DISEASE
- (iii) NUMBER OF SEQUENCES: 11
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Abbott Laboratories, D377/AP6D
  - (B) STREET: 100 Abbott Park Road
  - (C) CITY: Abbott Park
  - (D) STATE: IL
  - (E) COUNTRY: USA
  - (F) ZIP: 60064
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Diskette
  - (B) COMPUTER: IBM Compatible
  - (C) OPERATING SYSTEM: DOS
  - (D) SOFTWARE: FastSEQ Version 2.0
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 08/798692
  - (B) FILING DATE: 12-FEB-1997
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Becker, Cheryl L.
  - (B) REGISTRATION NUMBER: 35,441
  - (C) REFERENCE/DOCKET NUMBER: 6048.US.P1
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 847-935-1729
  - (B) TELEFAX: 847-938-2623
  - (C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1236 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGGCCGCCC GTCGGAGCCA GAAGCGGAGG GGGCGCCGGG GGGAGCCGGG  
 CACCGCCCTG 60

CTGGTCCCCG TCGCGCTGGG CCTGGGCCTG GCGCTGGCCT GCCTCGGCCT  
 CCTGCTGGCC 120

GTGGTCAGTT TGGGGAGCCG GGCATCGCTG TCCGCCCAGG AGCCTGCCCCA  
 GGAGGAGCTG 180

GTGGCAGAGG AGGACCAGGA CCCGTCGGAA CTGAATCCCC AGACAGAAGA  
 AAGCCAGGAT 240

CCTGCGCCTT TCCTGAACCG ACTAGTTCGG CCTCGAAGAA GTGCACCTAA  
 AGGCCGAAA 300

ACACGGGCTC GAAGAGCGAT CGCAGCCCAT TATGAAGTTC ATCCACGACC  
 TGGACAGGAC 360

GGAGCGCAGG CAGGTGTGGA CGGGACAGTG AGTGGCTGGG AGGAAGCCAG  
 AATCAACAGC 420

TCCAGCCCTC TGCCTACAA CCGCCAGATC GGGGAGTTTA TAGTCACCCG  
 GGCTGGGCTC 480

TACTACCTGT ACTGTCAGGT GCACTTTGAT GAGGGGAAGG CTGTCTACCT  
 GAAGCTGGAC 540

TTGCTGGTGG ATGGTGTGCT GGCCCTGCGC TGCCTGGAGG AATTCTCAGC  
 CACTGCGGCG 600

AGTTCCTCG GGCCCCAGCT CCGCCTCTGC CAGGTGTCTG GGCTGTTGGC  
 CCTGCGGCCA 660

GGGTCCTCCC TGCGGATCCG CACCCTCCCC TGGGCCCATC TCAAGGCTGC  
 CCCCTTCCTC 720

ACCTACTTCG GACTCTTCCA GGTTCACTGA GGGGCCCTGG TCTCCCCGCA  
 GTCGTCCCAG 780

GCTGCCGGCT CCCCTCGACA GCTCTCTGGG CACCCGGTCC CCTCTGCCCC  
 ACCCTCAGCC 840

GCTCTTTGCT CCAGACCTGC CCCTCCCTCT AGAGGCTGCC TGGGCCTGTT  
 CACGTGTTTT 900

CCATCCCACA TAAATACAGT ATTCCCACTC TTATCTTACA ACAACCCAC  
 CGCCCACTCT 960

CCACCTCACT AGCTCCCCAA TCCCTGACCC TTTGAGGCCC CCAGTGATCT  
 CGACTCCCC 1020

CTGGCCACAG ACCCCAGGG CATTGTGTTC ACTGTACTCT GTGGGCAAGG  
 ATGGGTCCAG 1080

GAATTCCTG TGGATTTTA AAACAGATAT TATTTT  
1236

(A) LENGTH: 249 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

[illegible]

Tyr Tyr Leu Tyr Cys Gln Val His Phe Asp Glu Gly Lys Ala Val  
 Tyr  
 165 170 175  
 Leu Lys Leu Asp Leu Leu Val Asp Gly Val Leu Ala Leu Arg Cys  
 Leu  
 180 185 190  
 Glu Glu Phe Ser Ala Thr Ala Ala Ser Ser Leu Gly Pro Gln Leu  
 Arg  
 195 200 205  
 Leu Cys Gln Val Ser Gly Leu Leu Ala Leu Arg Pro Gly Ser Ser  
 Leu  
 210 215 220  
 Arg Ile Arg Thr Leu Pro Trp Ala His Leu Lys Ala Ala Pro Phe  
 Leu  
 225 230 235  
 240  
 Thr Tyr Phe Gly Leu Phe Gln Val His  
 245

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 189 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: None

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Val Met Met Leu Arg Thr Trp Arg Leu Leu Pro Met Val Leu  
 Leu  
 1 5 10 15  
 Ala Ala Tyr Cys Tyr Cys Ser Leu Ala Ala Pro Gly Ser Asp Tyr  
 Lys  
 20 25 30  
 Asp Asp Asp Asp Lys Gly Arg Lys Thr Arg Ala Arg Arg Gly Ile  
 Ala  
 35 40 45  
 Ala His Tyr Glu Val His Pro Arg Pro Gly Gln Asp Gly Ala Gln  
 Ala  
 50 55 60  
 Gly Val Asp Gly Thr Val Ser Gly Trp Glu Glu Ala Arg Ile Asn  
 Ser  
 65 70 75  
 80  
 Ser Ser Pro Leu Arg Tyr Asn Arg Gln Ile Gly Glu Phe Ile Val  
 Thr  
 85 90 95  
 Arg Ala Gly Leu Tyr Tyr Leu Tyr Cys Gln Val His Phe Asp Glu  
 Gly  
 100 105 110  
 Lys Ala Val Tyr Leu Lys Leu Asp Leu Leu Val Asp Gly Val Leu  
 Ala

		115					120					125			
Leu	Arg	Cys	Leu	Glu	Glu	Phe	Ser	Ala	Thr	Ala	Ala	Ser	Ser	Leu	
Gly															
	130					135					140				
Pro	Gln	Leu	Arg	Leu	Cys	Gln	Val	Ser	Gly	Leu	Leu	Ala	Leu	Arg	
Pro															
145					150					155					
160															
Gly	Ser	Ser	Leu	Arg	Ile	Arg	Thr	Leu	Pro	Trp	Ala	His	Leu	Lys	
Ala															
				165					170						175
Ala	Pro	Phe	Leu	Thr	Tyr	Phe	Gly	Leu	Phe	Gln	Val	His			
			180					185							

(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met  Lys  Trp  Val  Thr  Phe  Ile  Ser  Leu  Leu  Phe  Leu  Phe  Ser  Ser
Ala
  1              5              10              15
  Tyr  Ser  Arg  Gly  Val  Phe  Arg  Arg
                20

```

(A) LENGTH: 8 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Tyr Lys Asp Asp Asp Asp Lys  
1 5

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 146 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Arg Arg Ala Ile Ala Ala His Tyr Glu Val His Pro Arg Pro Gly
Gln
1           5           10           15
Asp Gly Ala Gln Ala Gly Val Asp Gly Thr Val Ser Gly Trp Glu
Glu
          20           25           30
Ala Arg Ile Asn Ser Ser Ser Pro Leu Arg Tyr Asn Arg Gln Ile
Gly
          35           40           45
Glu Phe Ile Val Thr Arg Ala Gly Leu Tyr Tyr Leu Tyr Cys Gln
Val
        50           55           60
His Phe Asp Glu Gly Lys Ala Val Tyr Leu Lys Leu Asp Leu Leu
Val
        65           70           75
80
Asp Gly Val Leu Ala Leu Arg Cys Leu Glu Glu Phe Ser Ala Thr
Ala
          85           90           95
Ala Ser Ser Leu Gly Pro Gln Leu Arg Leu Cys Gln Val Ser Gly
Leu
        100           105           110
Leu Ala Leu Arg Pro Gly Ser Ser Leu Arg Ile Arg Thr Leu Pro
Trp
        115           120           125
Ala His Leu Lys Ala Ala Pro Phe Leu Thr Tyr Phe Gly Leu Phe
Gln
        130           135           140
Val His
145

```

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 177 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:



Glu Lys Gln Gln Asn Ile Ser Pro Leu Val Arg Glu Arg Gly Pro  
 Gln  
 1 5 10 15  
 Arg Val Ala Ala His Ile Thr Gly Thr Arg Gly Arg Ser Asn Thr  
 Leu  
 20 25 30  
 Ser Ser Pro Asn Ser Lys Asn Glu Lys Ala Leu Gly Arg Lys Ile  
 Asn  
 35 40 45  
 Ser Trp Glu Ser Ser Arg Ser Gly His Ser Phe Leu Ser Asn Leu  
 His  
 50 55 60  
 Leu Arg Asn Gly Glu Leu Val Ile His Glu Lys Gly Phe Tyr Tyr  
 Ile  
 65 70 75  
 80  
 Tyr Ser Gln Thr Tyr Phe Arg Phe Gln Glu Glu Ile Lys Glu Asn  
 Thr  
 85 90 95  
 Lys Asn Asp Lys Gln Met Val Gln Tyr Ile Tyr Lys Tyr Thr Ser  
 Tyr  
 100 105 110  
 Pro Asp Pro Ile Leu Leu Met Lys Ser Ala Arg Asn Ser Cys Trp  
 Ser  
 115 120 125  
 Lys Asp Ala Glu Tyr Gly Leu Tyr Ser Ile Tyr Gln Gly Gly Ile  
 Phe  
 130 135 140  
 Glu Leu Lys Glu Asn Asp Arg Ile Phe Val Ser Val Thr Asn Glu  
 His  
 145 150 155  
 160  
 Leu Ile Asp Met Asp His Glu Ala Ser Phe Phe Gly Ala Phe Ile  
 Val  
 165 170 175  
 Gly

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 183 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Glu Lys Gln Leu Ser Thr Pro Pro Leu Pro Arg Gly Gly Arg Pro  
 Gln  
 1 5 10 15  
 Lys Val Ala Ala His Ile Thr Gly Ile Thr Arg Arg Ser Asn Ser  
 Ala  
 20 25 30

[illegible][illegible][illegible]

- [illegible]

[illegible][illegible][illegible]

```

Val Leu Leu Thr His Thr Ile Ser Arg Ile Ala Val Ser Tyr Gln
Thr
65                      70                      75
80
Lys Val Asn Leu Leu Ser Ala Ile Lys Ser Pro Cys Gln Arg Glu
Thr
85                      90                      95
Pro Glu Gly Ala Glu Ala Lys Pro Trp Glu Pro Ile Tyr Leu Gly
Gly
100                    105                    110
Val Phe Gln Leu Glu Lys Gly Asp Arg Leu Ser Ala Glu Ile Asn
Arg
115                    120                    125
Pro Asp Tyr Leu Asp Phe Ala Glu Ser Gly Gln Val Tyr Phe Gly
Ile
130                    135                    140
Ile Ala Ile
145

```

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 145 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Leu Lys Pro Ala Ala His Leu Ile Gly Asp Pro Ser Lys Gln Asn
Ser
1          5          10          15
Leu Leu Trp Arg Ala Asn Thr Asp Arg Ala Phe Leu Gln Asp Gly
Phe
20          25          30
Ser Leu Ser Asn Asn Ser Leu Leu Val Pro Thr Ser Gly Ile Tyr
Phe
35          40          45
Val Tyr Ser Gln Val Val Phe Ser Gly Lys Ala Tyr Ser Pro Lys
Ala
50          55          60
Thr Ser Ser Pro Leu Tyr Leu Ala His Glu Val Gln Leu Phe Ser
Ser
65          70          75
80
Gln Tyr Pro Phe His Val Pro Leu Leu Ser Ser Gln Lys Met Val
Tyr
85          90          95
Pro Gly Leu Gln Glu Pro Trp Leu His Ser Met Tyr His Gly Ser
Ser
100         105         110
Gly Gln Leu Thr Gln Gly Asp Gln Leu Ser Thr His Thr Asp Gly
Ile
115         120         125

```

Ser Pro Arg Ser Glu  
165

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

[illegible]

## CLAIMS

1. A method of detecting the presence of target polynucleotide of TREPA in a test sample, comprising:

(a) contacting said test sample with at least one TREPA specific polynucleotide or complement thereof; and

(b) detecting the presence of said target polynucleotide of TREPA in the test sample, wherein said TREPA specific polynucleotide has at least 50% identity to polynucleotide SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

2. The method of claim 1 wherein said target polynucleotide of TREPA is attached to a solid phase prior to performing step (a).

3. A method for detecting mRNA of TREPA in a test sample, comprising:

(a) performing reverse transcription with at least one primer in order to produce cDNA;

(b) amplifying said cDNA obtained from step (a) by using other oligonucleotide primer(s) of TREPA as sense and antisense primer(s) in a first-stage amplification to obtain TREPA amplicon; and

(c) detecting the presence of said TREPA amplicon in the test sample, wherein said oligonucleotide primers of TREPA have at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

4. The method of claim 3 wherein said test sample is reacted with a solid phase prior to performing step (a) or step (b) or step (c).

5. The method of claim 3, wherein said detection step comprises utilizing a detectable label capable of generating a measurable signal.

10. A test kit useful for detecting TREPA polynucleotide in a test sample, comprising a container containing at least one TREPA polynucleotide having at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

11. A purified polynucleotide or fragment thereof derived from TREPA gene wherein said purified polynucleotide is capable of selectively hybridizing to the nucleic acid of said TREPA gene, and wherein said purified polynucleotide has at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

12. The purified polynucleotide of claim 11 wherein said purified polynucleotide is produced by recombinant techniques.

13. The purified polynucleotide of claim 12 wherein said polynucleotide produced by recombinant techniques comprises a sequence of at least one epitope encoded by TREPA.

14. A recombinant expression system comprising a nucleic acid sequence that encodes an open reading frame derived from TREPA which is operably linked to a control sequence compatible with a desired host and wherein said nucleic acid sequence has at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

15. The recombinant expression system of claim 14 further comprising a cell transformed with said recombinant expression system.

16. A polypeptide encoded by TREPA, wherein said polypeptide has at least 35% identity to amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

17. The polypeptide of claim 16 wherein said polypeptide is produced by recombinant technology.

18. The polypeptide of claim 16 wherein said polypeptide is produced by synthetic techniques.

19. A compound which inhibits activation of the TREPA polypeptide of claim 16.

20. The polypeptide of claim 16, wherein said polypeptide is a soluble fragment of the TREPA protein and is capable of binding a receptor for TREPA.

21. A method for treating a patient having a need to induce activation of the TREPA polypeptide of claim 16, comprising administering to said patient a therapeutically effective amount of a compound which induces activation of the TREPA polypeptide of claim 16.

22. A method for determining whether a compound is an agonist or antagonist to TREPA protein, comprising:

- (a) contacting a cell having TREPA protein expressed on its surface with said compound and a receptor ligand;
- (b) determining whether a biological effect is produced from the interaction of said cell and said compound; and
- (c) determining whether said compound is an agonist or antagonist.

23. A method for determining whether a receptor binds to a TREPA ligand, comprising:

- (a) contacting a mammalian cell which expresses the TREPA ligand with a receptor;
- (b) detecting the presence of the receptor; and
- (c) determining whether the receptor binds to the TREPA ligand.

24. An antibody which specifically binds to at least one epitope encoded by TREPA, wherein said antibody is polyclonal or monoclonal and wherein said epitope comprises an amino acid sequence having at least 35% identity to an



amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

25. An assay kit for determining the presence of TREPA antigen or antibody in a test sample, comprising a container containing a TREPA polypeptide having at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

26. The assay kit of claim 25 wherein said polypeptide is attached to a solid phase.

27. An assay kit for determining the presence of TREPA antigen or antibody in a test sample, comprising a container containing an antibody which specifically binds to TREPA antigen, wherein said antigen comprises at least one epitope of TREPA having at least about 60% similarity to a sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

28. The kit of claim 27 wherein said antibody is attached to a solid phase.

29. A method for producing a polypeptide comprising at least one epitope of TREPA, which method comprises incubating host cells transformed with an expression vector, wherein said vector comprises a polynucleotide sequence encoding a polypeptide, which polypeptide comprises an amino acid sequence having at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

30. A method for detecting TREPA antigen in a test sample suspected of containing said TREPA antigen, comprising:

(a) contacting said test sample with an antibody or fragment thereof which specifically binds to at least one

epitope of TREPA antigen selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof, for a time and under conditions sufficient for the formation of antibody/antigen complexes; and

(b) detecting said complexes.

31. The method of claim 30 wherein said antibody is attached to a solid phase.

32. A method for detecting antibodies which bind to TREPA antigen in a test sample suspected of containing said antibodies, comprising:

(a) contacting said test sample with a TREPA polypeptide, wherein said TREPA polypeptide contains at least one TREPA epitope comprising an amino acid sequence or fragment thereof having at least 35% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof, for a time and under conditions sufficient to allow antigen/antibody complexes to form;

(b) detecting said complexes.

33. The method of claim 32 wherein said TREPA polypeptide is attached to a solid phase.

34. A tissue culture grown cell comprising a nucleic acid sequence that encodes at least one epitope of TREPA antigen or a fragment thereof, wherein said nucleic acid sequence is transfected into said cell and wherein said nucleic acid sequence is SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

35. A method for producing antibodies which specifically bind to TREPA antigen, comprising administering to an individual an isolated immunogenic polypeptide or fragment thereof, wherein said isolated immunogenic polypeptide comprises at least one TREPA epitope and has at

least 35% identity to a sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof, in an amount sufficient to produce an immune response.

36. A method for producing antibodies which specifically bind to TREPA antigen, comprising administering to a mammal a plasmid comprising a sequence which encodes at least one epitope of TREPA, where said TREPA sequence is selected from the group consisting of SEQUENCE ID NO 1 and fragments or complements thereof.

37. A composition of matter comprising a TREPA polynucleotide or fragment thereof, wherein said polynucleotide has at least 50% identity to SEQUENCE ID NO 1 and fragments, analogs or complements thereof.

38. A composition of matter comprising a polypeptide containing at least one epitope encoded by TREPA, wherein said polypeptide has at least 35% identity to a sequence selected from the group consisting of SEQUENCE ID NO 2, SEQUENCE ID NO 3 and fragments thereof.

39. The composition of matter of claim 38, wherein said polypeptide is a soluble fragment of the TREPA protein and is capable of binding a receptor for TREPA.

40. The test kit of claim 10 further comprising a container containing tools useful for collection of said sample selected from the group consisting lancets, absorbent paper, cloth, swabs and cups.

41. The assay kit of claim 25 further comprising a container containing tools useful for collection of said sample selected from the group consisting lancets, absorbent paper, cloth, swabs and cups.

43. A gene or fragment thereof which codes for TREPA protein which comprises an amino acid sequence which has at least 35% identity to SEQUENCE ID NO 5.

44. A gene or fragment thereof comprising DNA having at least 35% identity to SEQUENCE ID NO 4.

**Abstract**—The purpose of this study was to determine the effect of a 10-week training program on the heart rate (HR) and energy expenditure (EE) of sedentary, middle-aged men. The subjects were divided into two groups: a control group (n = 10) and a training group (n = 10). The training group performed a 10-week program of aerobic exercise, consisting of 30 minutes of walking on a treadmill at 3.5 mph, 3 times per week. The control group did not exercise. HR and EE were measured at rest and during the exercise program. The training group showed a significant decrease in HR and an increase in EE compared to the control group. The results suggest that a 10-week training program can improve cardiovascular fitness and increase energy expenditure in sedentary, middle-aged men.

An isolated clone comprising sequences transcribed from the TREPA gene. Also provided are human polypeptides translated from said TREPA sequences and a procedure for producing such polypeptide by recombinant techniques. Also provided are a procedure for producing soluble biologically active TREPA, which may be used to treat deficiencies of TREPA and diseases conditions ameliorated by TREPA. Antibodies, antagonists and inhibitors of such polypeptide which may be used to prevent the action of such polypeptide and therefore may be used therapeutically to treat TREPA-associated diseases, tumors or metastases are disclosed. Also disclosed is the use of said antibodies, agonists and inhibitors as well as the nucleic acid sequences to screen for, diagnose, prognosticate, stage and monitor conditions and diseases attributable to TREPA, especially inflammation. The use of said partial sequence to provide antibodies, agonists and inhibitors as well as partial nucleic acid sequences to screen for, diagnose, stage and monitor diseases associated with TREPA, including but not limited to inflammation. Illustrative sequences and clone designations for TREPA are provided.

TREPA	104	- - - - - R R A I A A H Y E V h p r p g - - - - -	118
hTRAIL	105	E K Q Q N I S P L V R E R G P Q R V A A H I T G T R G R S N T L S S P N	140
mTRAIL	109	E K Q L S T P P L P R G G R P Q K V A A H I T G I T R R S N S A L I P I	144
hTNFa	86	- - - - - D K P V A H V V A N P Q A E - - - - -	99
hTNFb	61	- - - - - L K P A A H L I G D P S K Q - - - - -	74
h4-1BBL	89	- - - - - Q G M F A Q L V A g n v l i i d - - - - -	104
TREPA	119	- Q D G A Q A G v d g T V S G W E E A R - - - - I N S S S P L R Y N R Q	149
hTRAIL	141	S K N E K A L G R - - K I N S W E S S R - - - - S G H S F L S N L H L R	170
mTRAIL	145	S K D G K T L G Q - - K I E S W E S S R - - - - K G H S F L N H V L F R	174
hTNFa	100	- - - - - G Q L Q W L N R R - - - - A N A L L A N G V E L R	120
hTNFb	75	- - - - - N S L L W R A N T - - - - D R A F L Q D G F S L S	95
h4-1BBL	105	- - - - - G P L S W Y S D P g l a g V S L T G G L S Y K E D	129
TREPA	150	I G E F I V T R A G L Y Y L Y G Q V H F d e g k a v y i k l d l i v d g	185
hTRAIL	171	N G E L V I H E K G F Y Y I Y S Q T Y F r f q e e - - - - - I K	197
mTRAIL	175	N G E L V I E Q E G L Y Y I Y S Q T Y F r f q e a e d a s k m - - - V S	207
hTNFa	121	D N Q L V V P S E G L Y L I Y S Q V L F k g q g c p s t h v l i t h t i	156
hTNFb	96	N N S L L V P T S G I Y F V Y S Q V V F s g k a y s p k a t s s p i y i	131
h4-1BBL	130	T K E L V V A K A G V Y Y V F F Q L E L r r v v a g e g s g s v s - -	162
TREPA	186	v L A L R C L E E F S A T A A S S I g P Q L R L C Q V S G L L A L R P g	221
hTRAIL	198	E N T K N D K Q M V Q Y I Y K Y T S Y P D P I L L M K S A R N S C W s k	233
mTRAIL	208	K D K V R T K Q L V Q Y I Y K Y T S Y P D P I V L M K S A R N S C W s r	243
hTNFa	157	s r i a v s y q t k v n l i s a i k s p c q r e t p e g a e a k p w y e	192
hTNFb	132	a h e v q i f s - - - - - S Q Y P F H V P L L S S Q K m v y p g i	159
h4-1BBL	163	- L A L H L Q P L R S A A G A A I a l t v d i p p a s s e a r n s a -	196
TREPA	222	s s i - - - - R I R T L P W A H L K A A P F L - - - - -	240
hTRAIL	234	d a e y g i y S I Y Q G G I F E L K E N D R I f v s v t n e h l i d m d	269
mTRAIL	244	d a e y g i y S I Y Q G G L F E L K K N D R I f v s v t n e h l i d m d	279
hTNFa	193	- - - - - P I Y L G G V F Q L E K G D R L s a e i n r p d y l d f a	221
hTNFb	160	q e p w l h - S M Y H G A A F Q L T Q G D Q L s t h t d g i p h l v l s	194
h4-1BBL	197	- - - - - F G F Q G R L L H L S A G Q R L g v h l h t e a r a r h a	225
TREPA	241	- - - - - T Y F G L F q v h - - - - -	249
hTRAIL	270	h e a - - - - S F F G A F i v g - - - - -	281
mTRAIL	280	q e a - - - - S F F G A F i n - - - - -	291
hTNFa	222	e s g q - - - - V Y F G I i a l - - - - -	233
hTNFb	195	p s t - - - - V E F G A F a l - - - - -	205
h4-1BBL	226	w q l t q g a T V L G L F r v t P e i p a g l p s p r s e	254

Figure 1

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Steven R. Wiley

Continuation of U.S. Serial No.  
08/798,692, which was filed on  
February 12, 1997Title: MEMBER OF THE TNF FAMILY  
USEFUL FOR TREATMENT AND  
DIAGNOSIS OF DISEASE

Case No.: 6048.US.P1

Certificate of Mailing Under 37 C.F.R. §1.10

Express Mail Label No. EH 824 367 096 US

I hereby certify that this paper (along with any paper  
referred to as being attached or enclosed) is being  
deposited with the United States Postal Service on the  
date shown below as Express Mail Post Office to  
Addressee Service addressed to:Assistant Commissioner for Patents  
**Box Patent Application**  
Washington, D.C. 20231

Date of Deposit: February 10, 1998

Amy Martin

Date

DECLARATION AND POWER OF ATTORNEY FOR A UNITED STATES PATENT APPLICATIONAssistant Commissioner for Patents  
**Box Patent Application**  
Washington, D.C. 20231

As a below-named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name. I believe that I am an original and first and joint inventor of the subject matter of the invention entitled **MEMBER OF THE TNF FAMILY USEFUL FOR TREATMENT AND DIAGNOSIS OF DISEASE** which is claimed and for which a patent is sought in the patent application attached hereto.

I hereby state that I have reviewed and understand the contents of the above-mentioned specification, including the claims.

I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

Claim to benefit of foreign application(s):

I hereby claim foreign priority benefits under 35 U.S.C. §119 for the following foreign application(s) for patent or inventor's certificate:

NONE

The following foreign applications for patent or inventor's certificate have a filing date earlier than the filing date of the application(s) identified above:

NONE

08/798,692-01

Claim to benefit of earlier U.S. application(s):

I hereby claim the benefit under 35 U.S.C. §120 of the following earlier-filed United States patent application(s):

NONE

Insofar as the subject matter of each of the claims of this/these application(s) is not disclosed in the prior U.S. applications in the manner required by 35 U.S.C. §112, first paragraph, I acknowledge a duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. which came into existence between the filing date(s) of the prior applications and the national or PCT filing date of this application.

I hereby appoint the following Attorneys and/or agents to prosecute this application and any continuation or divisional applications based hereon, and to transact all business in the Patent and Trademark Office connected therewith:

Mona Anand, Reg. No. 34,537  
Regina M. Anderson, Reg. No. 35,820  
Mark C. Bach, Reg. No. 34,766  
Cheryl L. Becker, Reg. No. 35,441  
Thomas D. Brainard, Reg. No. 32,459  
Dianne Casuto, Reg. P-40,943  
Steven R. Crowley, Reg. No. 31,604  
Andreas M. Danckers, Reg. No. 32,652  
Michael J. Dixon, Reg. No. 32,410  
Matthew R. Hooper, Reg. No. 31,108

James D. McNeil, Reg. No. 26,204  
Neal D. Marcus, Reg. 35,267  
Lawrence S. Pope, Reg. No. 26,791  
Priscilla E. Porembski, Reg. No. 33,207  
Nicholas A. Poulos, Reg. No. 30,209  
Gregory W. Steele, Reg. No. 33,796  
Michael J. Ward, Reg. No. 37,960  
David L. Weinstein, Reg. No. 28,128  
Steven F. Weinstock, Reg. No. 30,117  
Brian R. Woodworth, Reg. No. 33,137  
Frank Z. Yang, Reg. No. 35,417  
Paul D. Yasger, Reg. No. 37,477

Send Correspondence to:

Steven F. Weinstock  
Abbott Laboratories  
D-377/AP6D  
100 Abbott Park Road  
Abbott Park, Illinois 60064-3500

Direct telephone calls to:

CHERYL L. BECKER  
(847)935-1729



